

Utilising invasive cacti as processed livestock feed and as a means of control

HO de Waal, Professional Animal Scientist/Researcher affiliated with the University of the Free State, Bloemfontein.
dewaalho@ufs.ac.za dewaalho0@gmail.com

Why this initiative?

Scientists at the University of the Free State have made major advances in utilising spineless cactus pears (*Opuntia ficus-indica* and *O. robusta*) as multi-use crops for humans and livestock. In addition to the conventional uses of the fruits, cladodes and fruits have been processed as livestock feed (De Waal 2015).

Construction of a wind turbine farm was started by an Independent Power Producer (IPP) near Bedford in the Eastern Cape Province. However, the construction footprint was heavily invaded by spiny cactus pears, notably the spiny form of *O. ficus-indica* and the spiny *O. engelmannii*. Acting on good advice the IPP called on the local expertise and knowledge in dealing with cactus pears, albeit the spineless forms.


The origin and extent of spiny cactus pear invasions

Invasive alien plants (IAPs) pose a direct threat to South Africa's biological diversity, as well as to water security, the ecological functioning of natural systems and productive use of land. A range of methods are used to control IAP's, namely mechanical, chemical, biological and integrated control methods (Anonymous 2009). In this initiative the focus is on mechanical control.

The invading alien spiny cactus pears are well-known in South Africa (NEMBA 2014a,b). Indications are that they were introduced by seafarers to the Cape of Good Hope about 300 years ago and later transported to the interior of the subcontinent (Kiesling & Metzger 2017). Spiny cactus pears resulted in dense, impenetrable thickets in some regions, especially the Eastern Cape Province (see **map**). By the 1950's about 1 million ha of South Africa had been invaded by the alien cacti (Anneck & Moran 1978 cited by Moran *et al.* 2013). Infestations of the spiny form of *O. ficus-indica*, so-called sweet prickly pear, severely impacted on agriculture (Moran *et al.* 2013), despite it being utilised by people and livestock.

In the early 1900's spineless cactus pear cultivars (*O. ficus-indica* and *O. robusta*) were imported from the Burbank collection in California to the Grootfontein Agricultural College in the Eastern Cape Province (De Kock 2001; Mondragón-Jacobo & Pérez-González 2001). During the past few decades, fruit production by spineless cactus pear cultivars (*O. ficus-indica*) for local markets and export has gained momentum. The plants are pruned annually to stimulate production of quality fruit and most of the fresh cladodes are used as livestock feed (De Waal 2015).

South Africa has a long history of living with and combating invasive alien plant species, including alien spiny cactus pear species (Moran *et al.* 2013). Spiny cactus pears have invaded large parts of the Eastern Cape Province and the spiny form of *O. ficus-indica* and another spiny cactus pear species *O. engelmannii* are of particular interest in this initiative. Although biological control agents have been introduced (**photo 1**), the success in controlling the invading alien cactus pear species varies (Zimmermann 2009).



Map. The distributions of sweet prickly pear (*Opuntia ficus-indica*) (black dots) and small round-leaved prickly pear (*O. engelmannii*) (red squares). Source: SAPIA database, ARC-PPRI.



Biological control of *O. engelmannii* using cochineal insects (Photo: HO de Waal)

Utilising invasive cacti as a means of control

Harvesting and processing spiny cactus pears

In addition to the continued actions of two biological control agents, mechanical control is used to harvest alien spiny cactus pears and the large volume of material is processed as livestock feed. Infestations will be opened, reclaimed, rehabilitated and the natural pastures (veld) allowed to revert back to grazing for livestock.

The harvesting and processing of the spiny cactus pears is fairly simple, although it requires a good measure of physical strength, perseverance and the necessary protective clothing. There are three successive stages, namely:

- harvesting of plants (**photo 2**),
- shredding through a cladode cutter (**photo 3a**) and drying in the sun (**photo 3b**),
- grinding in a hammer mill (**photo 4**).

The long spines are degraded mechanically by grinding the sun-dried cladode strips in a hammer mill before including it in balanced livestock diets.

Recommendations and the way forward

Cactus pears propagate vegetatively by means of cladodes sprouting roots and sexually through seed production. Over many centuries the spiny cactus pear fruits have been eaten by avian and mammal species, including people (without appropriate toilet facilities to contain the seeds); hence a vast seedbed has been created, just waiting for the right conditions to germinate.

During the cactus pear fruit ripening period, ripe fruits with mature seeds must be separated before harvesting and processing the cladodes. Although cladodes sprout roots and propagate vegetatively, dispersion of spiny cactus pear seeds in the faeces of animals (**photo 5**) and humans is a major means of spreading these invaders.

Mechanical control of alien spiny cactus pears is viewed as a long-term management activity, spanning a period of at least 20 years or more (Zimmermann *et al.* 2004). A range of methods is used to control IAP's, including mechanical, chemical, biological and integrated control methods. A control programme must include the following three phases:

- Initial control to drastically reduce the existing population.
- Follow-up control of seedlings, root suckers and coppice growth.
- Maintenance control on an annual basis to sustain low alien plant numbers.

The logistical challenges of rough terrain and the distances from the cacti harvesting sites to a processing facility are impacting conventional norms for profitable business planning. The National Resource Management Programme (NRMP) will be approached for official support in clearing and rehabilitating massive areas of infestations by alien spiny cactus pears in the Eastern Cape Province. It is envisaged that such financial support may be provided similar to that for the Working for Water (WfW) Programme (Zimmermann *et al.* 2004; Anonymous 2009; Moran *et al.* 2013) and will be an investment in reducing the infestations by invaders and the rehabilitation and sustainable use of natural resources in South Africa.



2



3a



3b



4

Photos 1—4: Francois Havenga



5

Photo 5: H O de Waal

Utilising invasive cacti as a means of control


The benefits

This initiative (starting in the Eastern Cape Province) is setting a baseline standard and creates an opportunity to manage the massive infestations of alien spiny cactus pears with mechanical control and to revert valuable under-utilised farm land back to natural grazing for livestock.

A primary objective of this baseline standard is to apply appropriate processing technology and transform the spiny plant material (through shredding, sun-drying and coarsely grinding in a hammer mill) into livestock feed which greatly limits the possibility of vegetative propagation.

An important spin-off and long-term benefit will also be gained from the large but untapped resource, namely converting invaders into useful processed livestock feed. Another important benefit is the range of employment opportunities which will be created for unemployed local people to engage in harvesting and processing the alien spiny cactus pears.

Photo 6: Shredded, sun-dried and coarsely ground spiny cactus pears (*Opuntia ficus-indica* and *O. engelmannii*), ready to be included in balanced diets for ruminant livestock (cattle, sheep, goats) and wild antelopes (photo: HO de Waal)



Literature consulted

Anneck, D.P. & Moran, V.C. 1978. Critical reviews of botanical pest control in South Africa 2: The prickly pear, *Opuntia ficus-indica* (L.) Miller. *J. Entomol. Soc. S. Afr.* 41, 161–188.

Anonymous. 2009. The war on invasive plants. *Wildlife Ranching* Winter 2009. Vol 2(3), 58–59.

De Kock, G.C. 2001. The use of *Opuntia* as a fodder source in arid areas of southern Africa. In: *FAO Plant Production and Protection Paper 169*. (eds.) Mondragón-Jacobo, C. & Pérez-González, S. Food and Agriculture Organization of the United Nations, Rome. pp 101–105.

De Waal, H.O. 2015. Spineless cactus pear as livestock feed in South Africa. In: *Development of a cactus pear agro-industry for the sub-Saharan Africa Region. Proceedings of International Workshop. 27–28 January 2015*. University of the Free State, Bloemfontein, South Africa. pp 30–39. *Cactusnet Newsletter*, 14 [Special Issue]. 94 pp.

Kiesling, R. & Metzger, D. 2017. Origin and taxonomy of *Opuntia ficus-indica*. In: Inglese P., Mondragon, C., Nefzaoui, A. & Saenz, C. (eds). *Crop ecology, cultivation and uses of cactus pear*. Advance draft prepared for the IX International Congress on Cactus Pear and Cochineal CAM crops for a hotter and drier world. Coquimbo, Chile, 26–30 March 2017. pp 29–35.

Mondragón-Jacobo, C. & Pérez-González, S. 2001. Germplasm resources and breeding *Opuntia* for fodder production. In: *FAO Plant Production and Protection Paper 169*. (eds.) Mondragón-Jacobo, C. & Pérez-González, S. Food and Agriculture Organization of the United Nations, Rome. pp 21–27.

Moran, V.C., Hoffmann, J.H. & Zimmermann, H.G. 2013. 100 years of biological control of invasive alien plants in South Africa: History, practice and achievements. *S. Afr. J. Sci.* 2013;109(9/10), Art. #a0022, 6 pages. <http://dx.doi.org/10.1590/sajs.2013/a0022>

NEMBA, 2014a. No. 598. National Environmental Management: Biodiversity Act 2004 (Act No. 10 of 2004): Alien and invasive Species Regulations, 2014. *Government Gazette*, 1 August 2014.

NEMBA, 2014b. No. 599. National Environmental Management: Biodiversity Act 2004 (Act No. 10 of 2004): Alien and invasive Species Lists, 2014. *Government Gazette*, 1 August 2014.

Zimmermann, H.G. 2009. Managing prickly pear invasions in South Africa. In: Nefzaoui, A., Inglese, P., Belay, T. (Eds.). *Improved utilization of cactus pear for food, feed, oil and water conservation and other products in Africa*. Proceedings of International Workshop, Mekelle (Ethiopia), 19–21 October, 2009, pp 157–166. *Cactusnet Newsletter*, 12 [Special Issue]. 224 pp.

Zimmermann, H.G., Moran, V.C. & Hoffmann, J.H. 2004. Biological control in the management of invasive alien plants in South Africa, and the role of the Working for Water programme. *S. Afr. J. Sci.* 100, 34–40.