

REFERENCES FOR BIOLOGICAL CONTROL TABLE UPDATED JULY 2018

- ADAIR, R.J. 2004. Seed-reducing Cecidomyiidae as potential biological control agents for invasive Australian wattles in South Africa, particularly *Acacia mearnsii* and *A. cyclops*. PhD Thesis, University of Cape Town, South Africa.
- ADAIR, R.J. 2005. The biology of *Dasineura dielsi* (Rübsaamen) (Diptera: Cecidomyiidae) in relation to the biological control of *Acacia cyclops* (Mimosaceae) in South Africa. *Australian Journal of Entomology* 44: 446-456.
- ADAIR, R.J., BURGESS, T., SERDANI, M. & BARBER, P. 2009. Fungal associations in *Asphondylia* (Diptera: Cecidomyiidae) galls from Australia and South Africa: implications for biological control of invasive acacias. *Fungal Ecology* 2: 121-134.
- ADAMSKI, D. & BROWN, R.L. 2002. A new species of *Frumenta* on *Solanum elaeagnifolium* in Mexico: a potential biocontrol agent for this noxious weed in southern Africa (Lepidoptera: Gelechiidae: Gnorimoschemini). *Proceedings of the Entomological Society of Washington* 104(4): 1029–1035.
- ANDERSON, R.S. 2015. *Pereskiophaga brasiliensis* Anderson, a new genus and species of Cryptorhynchinae (Coleoptera: Curculionidae) associated with primitive cacti of the genus *Pereskia* Miller (Cactaceae) in Brazil. *The Coleopterists Bulletin* 69(4): 670-673.
- ANNECKE, D.P. & MORAN, V.C. 1978. Critical reviews of biological pest control in South Africa. 2. The prickly pear, *Opuntia ficus-indica* (L.) Miller. *Journal of the Entomological Society of southern Africa* 41: 161-188.
- ANON. 1999. Appendix. In: Olckers, T. & Hill, M. P. (Eds) Biological Control of Weeds in South Africa (1990-1999). *African Entomology Memoir* No. 1. 175-182.
- BAARS, J-R. 2001. Biology and laboratory culturing of the root-feeding flea beetle, *Longitarsus columbicus columbicus* Harold, 1876 (Chrysomelidae: Alticinae): a potential natural enemy of *Lantana camara* L. (Verbenaceae) in South Africa. *Entomotropica* 16: 149-155.
- BAARS, J-R. 2002. The life history and host specificity of *Teleonemia vulgata* (Hemiptera: Tingidae), a potential biocontrol agent of *Lantana camara* (Verbenaceae). *African Entomology* 10: 315-324.
- BAARS, J-R. 2003. Geographic range, impact, and parasitism of lepidopteran species associated with the invasive weed *Lantana camara* in South Africa. *Biological Control* 28: 293-301.
- BAARS, J-R. & NESER, S. 1999. Past and present initiatives on the biological control of *Lantana camara* (Verbenaceae) in South Africa. In: Olckers, T. & Hill, M. P. (Eds) Biological Control of Weeds in South Africa (1990-1999). *African Entomology Memoir* No. 1. 21-33.
- BAARS, J-R., HILL, M.P., HEYSTEK, F., NESER, S. & URBAN, A.J. 2006. Biology, oviposition preference and impact in quarantine of the petiole-galling weevil, *Coelocephalapion camarae* Kissinger, a promising candidate agent for the biological control of *Lantana camara*. *Biological Control* 40: 187-195.
- BAARS, J-R., URBAN, A.J. & HILL, M.P. 2003. Biology, host range, and risk assessment supporting release in Africa of *Falconia intermedia* (Heteroptera: Miridae), a new biocontrol agent for *Lantana camara*. *Biological Control* 28: 282-292.

BALCIUNAS, J.K., BURROWS, D.W. & PURCELL, M.F. 1996. Comparison of the physiological and realized host-ranges of a biological control agent from Australia for the control of the aquatic weed, *Hydrilla verticillata*. *Biological Control* 7: 148-158.

BOWNES, A. 2009. Evaluation of a plant-herbivore system in determining potential efficacy of a candidate biological control agent, *Cornops aquaticum*, for water hyacinth, *Eichhornia crassipes*. PhD thesis, Rhodes University, Grahamstown.

BOWNES, A. 2010. Asian aquatic plant moth *Parapoynx diminutalis*, accidentally introduced earlier, contributes to control of an aquatic weed *Hydrilla verticillata* in South Africa. *African Journal of Aquatic Science* 35: 307-311.

BOWNES A. 2014. Suitability of a leaf-mining fly, *Hydrellia* sp., for biological control of the invasive aquatic weed, *Hydrilla verticillata* in South Africa. *BioControl* 59: 771-780.

BOWNES, A. 2015. A comparison of host range and performance of congeneric leaf-mining flies, *Hydrellia pakistanae* (Diptera: Ephydriidae) and *Hydrellia* sp., two candidate biological control agents for South African biotype of *Hydrilla verticillata* (Hydrocharitaceae). *Biological Control* 84: 44-52.

BOWNES, A. 2018. Suppression of the aquatic weed, *Hydrilla verticillata* (L.f.) Royle (Hydrocharitaceae) by a leaf-cutting moth *Parapoynx diminutalis* Snellen (Lepidoptera: Crambidae) in Jozini Dam, South Africa. *African Journal of Aquatic Science* 43: 153-162.

BOWNES, A., HILL, M.P. & BYRNE, M.J. 2013a. Nutrient-mediated effects on *Cornops aquaticum* Brünner (Orthoptera: Acrididae), a potential biological control agent of water hyacinth, *Eichhornia crassipes* (Mart.) Solms (Pontederiaceae). *Biological Control* 67: 548-554.

BOWNES, A., HILL, M.P. & BYRNE, M.J. 2013b. The role of nutrients in the responses of water hyacinth, *Eichhornia crassipes* (Pontederiaceae) to herbivory by a grasshopper *Cornops aquaticum* Brünner (Orthoptera: Acrididae). *Biological Control* 67: 555-562.

BOWNES, A. & DEEMING, J.C. 2016. A new species of *Hydrellia* Robineau-Desvoidy (Diptera: Ephydriidae) mining *Hydrilla verticillata* (L.f.) Royle (Hydrocharitaceae) leaves in Singapore. *Austral Entomology* 55: 353-359.

BOWNES, A., HILL, M.P. & BYRNE, M.J. 2010a. Evaluating the impact of herbivory by a grasshopper, *Cornops aquaticum* (Orthoptera: Acrididae), on the competitive performance and biomass accumulation of water hyacinth, *Eichhornia crassipes* (Pontederiaceae). *Biological Control* 53: 297–303.

BOWNES, A., HILL, M.P. & BYRNE, M.J. 2010b. Assessing density-damage relationships between water hyacinth and its grasshopper herbivore. *Entomologia Experimentalis et Applicata* 137: 246-254.

BROWN, J.W. 2006. A new species of *Cochylis* (Lepidoptera: Tortricidae: Cochylini) from Argentina: A Potential biocontrol agent against pompon weed, *Proceedings of the Entomological Society of Washington* 108 (4): 899-904

- BROWN, J.W. & ZACHARIADES, C. 2007. A new species of *Dichrorampha* (Lepidoptera: Tortricidae: Grapholitini) from Jamaica: a potential biocontrol agent against *Chromolaena odorata* (Asteraceae). *Proceedings of the Entomological Society of Washington* 109: 938-947.
- BUCCELLATO L., BYRNE, M.J. & WITKOWSKI, E.T.F. 2012. Interactions between a stem gall fly and a leaf-spot pathogen in the biological control of *Ageratina adenophora*. *Biological Control* 61: 222–229.
- BUCKINGHAM, G.R. & BALCIUNAS, J.K. 1994. Biological studies of *Bagous hydrillae*. US Army Corps of Engineers, Technical Report A-94-6.
- BUCKINGHAM, G.R., OKRAH, E.A. & THOMAS, M.C. 1989. Laboratory host range tests with *Hydrellia pakistanae* (Diptera: Ephydriidae), an agent for biological control of *Hydrilla verticillata* (Hydrocharitaceae). *Environmental Entomology* 18: 164-171.
- BYRNE, M.J., CURRIN, S. & HILL, M.P. 2002. The influence of climate on the establishment and success of the biocontrol agent, *Gratiana spadicea* released on *Solanum sisymbriifolium* in South Africa. *Biological Control* 24: 128-134.
- BYRNE, M.J., WITKOWSKI, E.T.F. & KALIBBALA, F.N. 2011. A review of recent efforts at biological control of *Caesalpinia decapetala* (Roth) Alston (Fabaceae) in South Africa. *African Entomology* 19: 247-257.
- CAGNOTTI, C., Mc KAY, F. & GANDOLFO, D. 2007. Biology and host specificity of *Plectonocha correntina* Lacordaire (Chrysomelidae), a candidate for the biological control of *Anredera cordifolia* (Tenore) Steenis. *African Entomology* 15: 300-309.
- CALDWELL, P.M. & KLUGE, R.L. 1993. Failure of the introduction of *Actinote antea* (Lep.: Acraeidae) from Costa Rica as a biological control candidate for *Chromolaena odorata* (Asteraceae) in South Africa. *Entomophaga* 38: 475-478.
- CENTER, T.D. & HILL, M.P. 2002. Field efficacy and predicted host range of the pickerelweed borer, *Bellura densa*, a potential biological control agent of water hyacinth. *BioControl* 47: 231-243.
- CENTER, T.D., GRODOWITZ, M.J., COFRANCESCO, A.F., JUKINSKY, G., SNODDY, E. & FREEDMAN, J.E. 1997. Establishment of *Hydrellia pakistanae* (Diptera: Ephydriidae) for the biological control of the submersed aquatic plant *Hydrilla verticillata* (Hydrocharitaceae) in the southeastern United States. *Biological Control* 8: 65-73.
- CHIDAWANYIKA, F., NYAMUKONDIWA, C., STRATHIE, L. & FISCHER, K. 2017. Effects of thermal regimes, starvation and age on heat tolerance of the parthenium beetle *Zygogramma bicolorata* (Coleoptera: Chrysomelidae) following dynamic and static protocols. *PLoS ONE* 12(1): e0169371. doi:10.1371/journal.pone.0169371
- CILLIERS, C.J. 1987. Notes on the biology of the established insect natural enemies of *Lantana camara* L. (Verbenaceae) and their seasonal history in South Africa. *Journal of the Entomological Society of Southern Africa* 46: 131-138.
- CILLIERS, C.J. 1991. Biological control of *Pistia stratiotes* (Araceae) in South Africa. *Agriculture, Ecosystems and Environment* 37: 225-229.

CILLIERS, C.J. 1999. Biological control of parrot's feather, *Myriophyllum aquaticum* (Vell.) Verdc. (Haloragaceae) in South Africa. In: Olckers, T. & Hill, M. P. (Eds) Biological Control of Weeds in South Africa (1990-1999). *African Entomology Memoir* No. 1. 113-118.

CILLIERS, C.J. & NESER, S. 1991. Biological control of *Lantana camara* (Verbenaceae) in South Africa. *Agriculture, Ecosystems and Environment* 37: 57-75.

CILLIERS, C.J., HILL, M.P., OGWANG, J.A. & AJUONU, O. 2003. Aquatic weeds in Africa and their control. In: Neuenschwander, P., Borgemeister, C. & Langewald, J. (Eds) *Biological Control in IPM Systems in Africa*. 161–178. CAB International, Wallingford.

COETZEE, J.A., BYRNE, M.J. & HILL, M.P. 2007. Impact of nutrients and herbivory by *Eccritotarsus catarinensis* on the biological control of water hyacinth, *Eichhornia crassipes*. *Aquatic Botany* 86: 179-186.

COETZEE, J.A., CENTER, T.D., BYRNE, M.J. & HILL, M.P. 2005. Impact of the biocontrol agent *Eccritotarsus catarinensis*, a sap-sucking mirid, on the competitive performance of water hyacinth, *Eichhornia crassipes*. *Biological Control* 32: 90-96.

COETZEE, J.A., HILL, M.P., JULIEN, M.H., CENTER, T.D. & CORDO, H.A. 2009. *Eichhornia crassipes*. In: Muniappan, R., Reddy, G.V.P. & Raman, A. (Eds.) *Biological Control of Tropical Weeds using Arthropods*. 183-210. Cambridge University Press, Cambridge.

COETZEE, J.A., BOWNES, A. & MARTIN, G.D. 2011a. Prospects for the biological control of submerged macrophytes in South Africa. *African Entomology* 19: 469-487

COETZEE, J.A., HILL, M.P., BYRNE, M.J. & BOWNES, A. 2011b. A review of the biological control programmes on *Eichhornia crassipes* (C.Mart.) Solms (Pontederiaceae), *Salvinia molesta* D.S.Mitch. (Salviniaceae), *Pistia stratiotes* L. (Araceae), *Myriophyllum aquaticum* (Vell.) Verdc. (Haloragaceae) and *Azolla filiculoides* Lam. (Azollaceae) in South Africa. *African Entomology* 19: 451-468.

COETZER, W. 2000. Oviposition preference of *Sulcobruchus subsuturalis* (Pic) (Coleoptera: Bruchidae), an introduced natural enemy of *Caesalpinia decapetala* (Roth) Alston (Caesalpinaceae) in South Africa. *African Entomology* 8: 293-297.

COETZER, W. & HOFFMANN, J.H. 1997. Establishment of *Neltumius arizonensis* (Coleoptera: Bruchidae) on mesquite (*Prosopis* species: Mimosaceae) in South Africa. *Biological Control* 10: 187-192.

COETZER, W. & NESER, S. 1999. Biological control initiatives against the invasive Oriental legume, *Caesalpinia decapetala* (Roth) Alston (Mauritius thorn), in South Africa. In: Olckers, T. & Hill, M. P. (Eds) Biological Control of Weeds in South Africa (1990-1999). *African Entomology Memoir* No. 1. 145-152.

CORDO, H.A. & DELOACH, C.J. 1982. Weevils *Listronotus marginicollis* and *L. cinnamomeus* that feed on *Limnobium* and *Myriophyllum* in Argentina. *The Coleopterists Bulletin* 36: 302-308.

COWIE, B.W., BYRNE, M.J., WITKOWSKI, E.T.F & VENTER, N. 2016b. Exacerbation of photosynthetic damage through increased heat–light stress resulting from *Gargaphia decoris* sap-feeding. *Biological Control* 94: 82-89

COWIE, B.W., VENTER, N., WITKOWSKI, E.T.F., BYRNE, M.J. & OLCKERS, T. 2018. A review of *Solanum mauritianum* biocontrol: prospects, promise and problems: a way forward for South Africa and globally. *BioControl* (DOI: 10.1007/s10526-017-9858-0)

COWIE, B.W., VENTURI, G., WITKOWSKI, E.T.F. & BYRNE, M.J. 2016a. Does climate constrain the spread of *Anthonomus santacruzi*, a biological control agent of *Solanum mauritianum*, in South Africa? *Biological Control* 101: 1–7.

COWIE, B.W., WITKOWSKI, E.T.F., BYRNE, M.J. & VENTER, N. 2017. A villainous hero: does the biological control agent, *Anthonomus santacruzi*, pollinate its target weed, *Solanum mauritianum*? *Biological Control* 105: 79–85.

CRAEMER, C. 1995. Host specificity, and release in South Africa, of *Aceria malherbae* Nuzzaci (Acari: Eriophyoidea), a natural enemy of *Convolvulus arvensis* L. (Convolvulaceae). *African Entomology* 3: 213-215.

CRAEMER, C., NESER, S. & SMITH MEYER, M.K.P. 1996. Eriophyid-myte (Acari: Eriophyoidea: Eriophyidae) as moontlike beheeragente van ongewenste uitheemse plante in Suid-Afrika. *Suid-Afrikaanse Tydskrif vir Natuurwetenskap en Tegnologie* 15: 99-109.

DAY, M.D., WILEY, C.J., PLAYFORD, J. & ZALUCKI, M.P. 2003. *Lantana: Current Management Status and Future Prospects*. ACIAR Monograph No.102. Australian Centre for International Agricultural Research, Canberra. [<http://www.aciar.gov.au/publication/MN102>]

DELGADO, O.S., STRATHIE, L.W. & ZACHARIADES, C. 2014. Host range of *Conotrachelus reticulatus*, a potential biological control agent for *Chromolaena odorata*. In: *Proceedings of the XIV International Symposium on Biological Control of Weeds*, pp 31-40. F.A.C. Impson, C.A. Kleinjan and J.H. Hoffmann (eds). 2-7 March 2014, Kruger National Park, South Africa. University of Cape Town, South Africa.

DELOACH, C.J., LEWIS, P.A., HERR, J.C., CARRUTHERS, R.I., TRACY, J.L. & JOHNSON, J. 2003. Host specificity of the leaf beetle, *Diorhabda elongata deserticola* (Coleoptera: Chrysomelidae) from Asia, a biological control agent for saltcedars (*Tamarix*: Tamaricaceae) in the Western United States. *Biological Control* 27: 117-147.

DEN BREEÏEN, A. 2003. Release strategies for the establishment of the leaf spot pathogen, *Mycovellosiella lantanae* var. *lantanae*, on *Lantana camara* in South Africa. In: Cullen, J.M., Briese, D.T., Kriticos, D.J., Lonsdale, W.M., Morin, L. & Scott, J.K. (Eds) *Proceedings of the XI International Symposium on Biological Control of Weeds*. 386-388. CSIRO Entomology, Canberra.

DEN BREEÏEN, A. & MORRIS, M.J. 2003. Pathogenicity and host specificity of *Mycovellosiella lantanae* var. *lantanae*, a potential biocontrol agent for *Lantana camara* in South Africa. *Biocontrol Science and Technology* 13: 313 – 322.

DEN BREEÏEN, A., GROENEWALD, J.Z., VERKLEY, G.J.M. & CROUS, P.W. 2006. Morphological and molecular characterisation of *Mycosphaerellaceae* associated with the invasive weed, *Chromolaena odorata*. *Fungal Diversity* 23: 89-110.

DENNILL, G.B. & DONNELLY, D. 1991. Biological control of *Acacia longifolia* and related weed species (Fabaceae) in South Africa. *Agriculture, Ecosystems and Environment* 37: 115-135.

- DENNILL, G.B., DONNELLY, D., STEWART, K., & IMPSON, F.A.C. 1999. Insect agents used for the biological control of Australian *Acacia* species and *Paraserianthes lophantha* (Willd.) Nielsen (Fabaceae) in South Africa. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology Memoir* No. 1. 45-54.
- DHILEEPAN, K. & STRATHIE, L. 2009. *Parthenium hysterophorus* L. (Asteraceae). In: Muniappan, R., Reddy, G.V.P. & Raman, A. (Eds) *Biological Control of Tropical Weeds using Arthropods*. 274-318. Cambridge University Press, Cambridge.
- DHILEEPAN, K., SNOW, E.L., RAFTER, M.A., TREVIÑO, M., MCCARTHY, J. & WILMOT SENARATNE, K.A.S. 2007. The leaf-tying moth *Hypocosmia pyrochroma* (Lep., Pyralidae), a host-specific biological control agent for cat's claw creeper *Macfadyena unguis-cati* (Bignoniaceae) in Australia. *Journal of Applied Entomology* 131: 564-568.
- DI IORIO, O.R. 1998. Cerambycidae (Coleoptera) root and stem borers in living plants from Argentina: a bibliographical review and new host plant records. *Giornale Italiano di Entomologia* 9: 73-95.
- DORCHIN, N. & ADAIR, R.J. 2011. Two new *Dasineura* species (Diptera: Cecidomyiidae) from coastal tea tree, *Leptospermum laevigatum* (Myrtaceae) in Australia. *Australian Journal of Entomology* 50: 65-71.
- DUBE, N., ASSEFA, Y., ZACHARIADES, C., CONLONG, D. & OLCKERS, T. 2014. Genetic diversity in *Pareuchaetes insulata* and its implications for biological control of *Chromolaena odorata*. *BioControl* 59: 253-262.
- DUBE, N., ZACHARIADES, C. & OLCKERS, T. 2009. Comparative biology and effects of crossbreeding *Pareuchaetes insulata* (Lepidoptera: Arctiidae), a biological control agent of *Chromolaena odorata* (Asteraceae). *Proceedings of the South African Sugar Technologists' Association* 82: 433 – 436.
- DUBE, N., ZACHARIADES, C., MUNYAI, T.C. & UYI, O.O. 2017. Laboratory studies on the biology and host range of *Dichrorampha odorata* (Lepidoptera: Tortricidae), a biological control agent for *Chromolaena odorata* (Asteraceae). *Biocontrol Science and Technology* 27 (2): 222–236.
- EICHLIN, T.D., DELGADO, O.S., STRATHIE, L.W., ZACHARIADES, C. & CLAVIJO, J. 2009. *Carmenta chromolaenae* Eichlin, a new species (Lepidoptera: Sesiidae) for the biological control of *Chromolaena odorata* (L.) King & Robinson (Asteraceae). *Zootaxa* 2288: 42-50.
- ENGLISH, K.F. & OLCKERS, T. 2018. Abundance across seasons and early impact of the florivorous weevil *Anthonomus santacruzii* (Curculionidae) on *Solanum mauritianum* (Solanaceae) in KwaZulu-Natal Province, South Africa. *Biological Control* 121: 175-182.
- ERASMUS, D.J., BENNETT, P.H. & VAN STADEN, J. 1992. The effect of galls induced by the gall fly *Procecidochares utilis* on vegetative growth and reproductive potential of crofton weed, *Ageratina adenophora*. *Annals of Applied Biology* 120: 173-181.
- FOURIE, A. 2011. Preliminary attempts to identify pathogens as biological control agents against *Cestrum* species (Solanaceae) in South Africa. *African Entomology* 19: 278-281.
- FOWLER, S.V., BARRETO, R., DODD, S., MACEDO D.M., PAYNTER, Q., PEDROSA-MACEDO, J.H., PEREIRA, O.L., PETERSON, P., SMITH, L., WAIPARA, N., WINKS, C.J. & FORRESTER, G.

2013. *Tradescantia fluminensis*, an exotic weed affecting native forest regeneration in New Zealand: ecological surveys, safety tests and releases of four biocontrol agents from Brazil. *Biological Control* 64: 323-329.

GOOLSBY, J.A. & MORAN, P. 2009. Host range of *Tetramesa romana* Walker (Hymenoptera: Eurytomidae), a potential biological control agent of giant reed, *Arundo donax* L. in North America. *Biological Control* 49: 160-168.

GOOLSBY, J.A., MORAN, P.J., ADAMCZYK, J.J., KIRK, A.A., JONES, W.A., MARCOS, M.A. & CORTÉS, E. 2010. Host range of the European, rhizome-stem feeding scale *Rhizaspidotus donacis* (Hemiptera: Diaspididae), a candidate biological control agent for giant reed, *Arundo donax* (Poales: Poaceae) in North America. *Biocontrol Science and Technology* 19: 899-918.

GORDON, A.J. 1999a. A review of established and new insect agents for the biological control of *Hakea sericea* Schrader (Proteaceae) in South Africa. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology Memoir No. 1*. 35-43.

GORDON, A.J. 1999b. Biological control of Australian myrtle, *Leptospermum laevigatum* (J. Gaertn.) F. Muell. (Myrtaceae), in South Africa. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology Memoir No. 1*. 139-143.

GORDON, A.J. 2003. Biology and host range of the stem-boring beetle *Aphanasium australe*, a promising agent for the biological control of *Hakea sericea* in South Africa. *BioControl* 48: 113-122.

GORDON, A.J. 2011. Biological control endeavours against Australian myrtle, *Leptospermum laevigatum* (Gaertn.) F. Muell. (Myrtaceae), in South Africa. *African Entomology* 19: 349-355.

GORDON, A.J. & FOURIE, A. 2011. Biological control of *Hakea sericea* Schrad. & J.C. Wendl. and *Hakea gibbosa* (Sm.) Cav. (Proteaceae) in South Africa. *African Entomology* 19: 303-314.

GORDON, A.J. & KLUGE, R.L. 1991. Biological control of St. John's Wort, *Hypericum perforatum* (Clusiaceae) in South Africa. *Agriculture, Ecosystems and Environment* 37: 77-90.

HERR, J.C., HERRERA-REDDY, A.M. & CARRUTHERS, R.I. 2014. Field testing *Diorhabda elongata* (Coleoptera: Chrysomelidae) from Crete, Greece, to assess potential impact on nontarget native California plants in the genus *Frankenia*. *Environmental Entomology* 43: 642-653.

HENRY, T.J. 2017. Description of a cryptic new species of the plant bug genus *Eccritotarsus* (Heteroptera: Miridae: Bryocorinae) from Peru, a new biocontrol agent of water hyacinth, *Eichhornia crassipes* (Pontederiaceae). *Proceedings of the Entomological Society of Washington* 119: 398-407.

HEYSTEK, F. & BAARS, J.-R. 2005. Biology and host range of *Aconophora compressa*, a candidate considered as a biological agent of *Lantana camara* in Africa. *BioControl* 50: 359-373.

HEYSTEK, F., WOOD, A.R., NESER, S. & KISTENSAMY, Y. 2011. Biological control of two *Ageratina* species (Asteraceae: Eupatorieae) in South Africa. *African Entomology* 19: 208-216.

HILL, M. P. 1999. *Gratiana lutescens* (Boheman) (Coleoptera: Chrysomelidae: Cassidinae) reconsidered as a natural enemy for satansbos (*Solanum elaeagnifolium* Canavilles) (Solanaceae) in South Africa. *African Entomology* 7: 177-181.

- HILL, M.P. & CILLIERS, C.J. 1999. A review of the arthropod natural enemies, and factors that influence their efficacy, in the biological control of water hyacinth, *Eichhornia crassipes* (Mart.) Solms-Laub. (Pontederiaceae), in South Africa. In: Olckers, T. & Hill, M. P. (Eds) Biological Control of Weeds in South Africa (1990-1999). *African Entomology* Memoir No. 1. 103-112.
- HILL, M.P. & HULLEY, P.E. 1995. Biology and host range of *Gratiana spadicea* (Klug, 1829) (Coleoptera: Chrysomelidae: Cassidinae), a potential biological control agent for the weed *Solanum sisymbriifolium* Lamarck (Solanaceae) in South Africa. *Biological Control* 5: 345-352.
- HILL, M.P. & HULLEY, P.E. 1996. Suitability of *Metriorhina elatior* (Klug) (Coleoptera: Chrysomelidae: Cassidinae) as a biological control agent for *Solanum sisymbriifolium* Lam. (Solanaceae). *African Entomology* 4: 117-123.
- HILL, M.P. & McCONNACHIE, A.J. 2009. *Azolla filiculoides*. In: Muniappan, R., Reddy, G.V.P. & Raman, A. (Eds.) *Biological Control of Tropical Weeds using Arthropods*. 74-87. Cambridge University Press, Cambridge.
- HILL, M.P. & OBERHOLZER, I.G. 2002. Laboratory host range testing of the flea beetle, *Pseudolampsis guttata* (LeConte) (Coleoptera: Chrysomelidae), a potential natural enemy for red water fern, *Azolla filiculoides* Lamarck (Pteridophyta: Azollaceae) in South Africa. *The Coleopterists Bulletin* 56: 79-83.
- HILL, M.P., McCONNACHIE, M.J. & BYRNE, M.J. 2008. *Azolla filiculoides* Lamarck (Pteridophyta: Azollaceae) control in South Africa: a 10 year review. In: Julien, M.H., Sforza, R., Bon, M.C., Evans, H.C., Hatcher, P.E., Hinz, H.L. & Rector, B.G. (Eds) *Proceedings of the XII International Symposium on Biological Control of Weeds*. 566-568. CAB International Wallingford, UK.
- HODSON, J.L., HOFFMANN, J.H. & ZIMMERMANN, H.G. 2003. Biological control of spear thistle, *Cirsium vulgare* (Asteraceae), in South Africa: a modest start for *Rhinocyllus conicus* (Coleoptera: Curculionidae). *African Entomology* 11: 15-20.
- HOFFMANN, J.H. & MORAN, V.C. 1991. Biological control of *Sesbania punicea* (Fabaceae) in South Africa. *Agriculture, Ecosystems and Environment* 37: 157-173.
- HOFFMANN, J.H. & MORAN, V.C. 1999. A review of the agents and factors that have contributed to the successful biological control of *Sesbania punicea* (Cav.) Benth. (Papilionaceae) in South Africa. In: Olckers, T. & Hill, M. P. (Eds) Biological Control of Weeds in South Africa (1990-1999). *African Entomology* Memoir No. 1. 75-79.
- HOFFMANN, J.H., IMPSON, F.A.C., MORAN, V.C. & DONNELLY, D. 2002. Biological control of invasive golden wattle trees (*Acacia pycnantha*) by a gall wasp, *Trichilogaster* sp. (Hymenoptera: Pteromalidae), in South Africa. *Biological Control* 25: 64-73.
- HOFFMANN, J.H., IMPSON, F.A.C. & MORAN, V.C. 1993. Competitive interactions between two bruchid species (*Algarobius* spp.) introduced into South Africa for biological control of mesquite weeds (*Prosopis* spp.). *Biological Control* 3: 215-220.
- HOFFMANN, J.H., MORAN, V.C. & IMPSON, F.A.C. 1998. Promising results from the first biological control programme against a solanaceous weed (*Solanum elaeagnifolium*). *Agriculture, Ecosystems and Environment* 70: 145-150.

- HOFFMANN, J.H., MORAN, V.C. & VAN WILGEN, B.W. 2011. Prospects for the biological control of invasive *Pinus* species (Pinaceae) in South Africa. *African Entomology* 19: 393-401.
- HOFFMANN, J.H., MORAN, V.C. & ZIMMERMANN, H.G. 1999. Integrated management of *Opuntia stricta* (Haworth) Haworth (Cactaceae) in South Africa: an enhanced role for two, renowned, insect agents. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology* Memoir No. 1. 15-20.
- IMPSON, F.A.C., HOFFMANN, J.H. & KLEINJAN, C. 2009. Australian *Acacia* species (Mimosaceae) in South Africa. In: Muniappan, R., Reddy, G.V.P. & Raman, A. (Eds) *Biological Control of Tropical Weeds using Arthropods*. 38-62. Cambridge University Press, Cambridge.
- IMPSON, F.A.C., MORAN, V.C. & HOFFMANN, J.H. 1999. A review of the effectiveness of seed-feeding bruchid beetles in the biological control of mesquite, *Prosopis* species (Fabaceae), in South Africa. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology* Memoir No. 1. 81-88. Entomological Society of Southern Africa, Pretoria.
- IMPSON, F.A.C., KLEINJAN, C.A., HOFFMANN, J.H., POST, J.A. & WOOD, A.R. 2011. Biological control of Australian *Acacia* species and *Paraserianthes lophantha* (Willd.) Nielsen (Mimosaceae) in South Africa. *African Entomology* 19: 186-207.
- IMPSON, F.A.C., KLEINJAN, C.A., HOFFMANN, J.H. & POST, J.A. 2008. *Dasineura rubiformis* (Diptera: Cecidomyiidae), a new biological control agent for *Acacia mearnsii* in South Africa. *South African Journal of Science* 104: 247-249.
- JULIEN, M.H., HILL, M.P. & TIPPING, P.W. 2009. 25. *Salvinia molesta*. In: Muniappan, R., Reddy, G.V.P. & Raman, A. (Eds.) *Biological Control of Tropical Weeds using Arthropods*. 378-407. Cambridge University Press, Cambridge.
- KING, A.M., BRUDVIG, R. & BYRNE, M.J. 2011a. Biological control of dense-thorned bitter apple, *Solanum sisymbriifolium* Lam. (Solanaceae), in South Africa. *African Entomology* 19: 427-433.
- KING, A.M., WILLIAMS, H.E. & MADIRE, L.G. 2011b. Biological control of cat's claw creeper, *Macfadyena unguis-cati* (L.) A.H.Gentry (Bignoniaceae), in South Africa. *African Entomology* 19: 366-377.
- KLEIN, H. 1999. Biological control of three cactaceous weeds, *Pereskia aculeata* Miller, *Harrisia martinii* (Labouret) Britton and *Cereus jamacaru* De Candolle in South Africa. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology* Memoir No. 1. 3-14.
- KLUGE, R.L. 1991. Biological control of crofton weed, *Ageratina adenophora* (Asteraceae), in South Africa. *Agriculture, Ecosystems and Environment* 37: 187-191.
- KLUGE, R.L. & CALDWELL, P.M. 1993a. The biology and host specificity of *Pareuchaetes aurata aurata* (Lepidoptera: Arctiidae), a "new association" biological control agent for *Chromolaena odorata* (Compositae). *Bulletin of Entomological Research* 83: 87-94.
- KLUGE, R.L. & CALDWELL, P.M. 1993b. Host specificity of *Pareuchaetes insulata* (Lep.: Arctiidae), a biological control agent for *Chromolaena odorata* (Compositae). *Entomophaga* 38: 451-457.

- KLUGE, R.L. & NESER, S. 1991. Biological control of *Hakea sericea* (Proteaceae) in South Africa. *Agriculture, Ecosystems and Environment* 37: 91-113.
- KLUGE, R.L. & ZACHARIADES, C. 2006. Assessing the damage potential of the stem-boring weevil *Lixus aemulus* for the biological control of *Chromolaena odorata*. *BioControl* 51: 547-552.
- LENNOX, C.L., HOFFMANN, J.H., COUTINHO, T.A. & ROQUES, A. 2009. A threat of exacerbating the spread of pitch canker precludes further consideration of a cone weevil, *Pissodes validirostris*, for biological control of invasive pines in South Africa. *Biological Control* 50: 179-184.
- KOTZÉ, L.J.D., WOOD, A.R. & LENNOX, C.L. 2015. Risk assessment of the *Acacia cyclops* dieback pathogen, *Pseudolagarobasidium acaciicola*, as a mycoherbicide in the South African strandveld and limestone fynbos. *Biological Control* 82: 52-60.
- LENNOX, C.L., SERDANI, M., GROENEWALD, J.Z. & CROUS, P.W. 2004. *Prospodicola mexicana* gen. et. sp. nov., causing a new pod disease of *Prosopis* species. *Studies in Mycology* 50: 187-194.
- LUBBE, C.M., DENMAN, S., CANNON, P.F., GROENEWALD, J.Z., LAMPRECHT, S.C. & CROUS, P.W. 2004. Characterization of *Colletotrichum* species associated with diseases of Proteaceae. *Mycologia* 96: 1268-1279.
- MABUDA, K. 2005. *Eutreta xanthochaeta* (Diptera: Tephritidae) is unsuitable for biocontrol of *Lantana camara* in Africa. *Biocontrol Science and Technology* 15: 635-639.
- MADEIRA, P.T., HILL, M.P., DRAY JR, F.A., COETZEE, J.A., PATERSON, I.D. & TIPPING, P.W. 2016. Molecular identification of *Azolla* invasions in Africa: the *Azolla* specialist, *Stenopelmus rufinusus* proves to be an excellent taxonomist. *South African Journal of Botany* 105: 299-305
- MADIRE, L.G. 2010. Suitability of the leaf-mining fly, *Pseudonapomyza* sp. (Diptera: Agromyzidae), for biological control of *Tecoma stans* L. (Bignoniaceae) in South Africa. MSc thesis. University of Fort Hare, Alice, South Africa.
- MADIRE, L.G. 2013. Biology and host range of *Mada polluta*, a potential biological control agent of *Tecoma stans* in South Africa. *Biocontrol Science and Technology* 23(8): 944-955.
- MADIRE, L.G., WOOD, A.R., WILLIAMS, H.E. & NESER, S. 2011a. Potential agents for the biological control of *Tecoma stans* (L.) Juss. ex Kunth var. *stans* (Bignoniaceae) in South Africa. *African Entomology* 19: 434-442.
- MADIRE, L.G., SIMELANE, D.O. & WALADDE, S. 2011b. Biology and host range of the leafminer, *Pseudonapomyzasp.* (Diptera: Agromyzidae), a potential biological control agent for *Tecoma stans* (Bignoniaceae) in South Africa. *Biocontrol Science and Technology* 21(12): 1409-1421.
- MARLIN, D., NEWETE, S.W., MAYONDE, S.G. SMIT, E.R., & BYRNE, M.J. 2017. Invasive *Tamarix* (Tamaricaceae) in South Africa: current research and the potential for biological control. *Biological Invasions* 19 (10): 2971–2992.
- MARSHALL, M., GOOLSBY, J.A., VACEK, A.T., MORAN, P.J., KIRK, A.A., CORTES MENDOZA, E., CRISTOFARO, M., BOWNES, A., MASTORAS, A., KASHEFI, J., CHASKOPOULOU, A., SMITH, L., GOLDSMITH, B. & RACELIS, A.E. 2018. Densities of the arundo wasp, *Tetramesa romana* (Hymenoptera: Eurytomidae) across its native range in Mediterranean Europe and introduced

ranges in North America and Africa. *Biocontrol Science and Technology* DOI: 10.1080/09583157.2018.1493090

MARTINS, U.J. & GALILEO, M.H.M. 2004. Novos taxons de Aerenicini (Coleoptera, Cerambycidae, Lamiinae) da Guatemala, Brasil e Bolivia. *Iheringia, Serie Zoologia* 94: 395-401.

MATHENGE, C.W., HOLFORD, P., HOFFMANN, J.H., SPOONER-HART, R., BEATTIE G.A.C., & ZIMMERMANN, H.G. 2009a. The biology of *Dactylopius tomentosus* (Lamarck) (Hemiptera: Dactylopiidae). *Bulletin of Entomological Research* 99: 551-559.

MATHENGE, C.W., HOLFORD, P., HOFFMANN, J.H., SPOONER-HART, R., BEATTIE, G.A.C. & ZIMMERMANN H.G. 2009b. Distinguishing suitable biotypes of *Dactylopius tomentosus* (Hemiptera: Dactylopiidae) for biological control of *Cylindropuntia fulgida* var. *fulgida* (Caryophyllales: Cactaceae) in South Africa. *Bulletin of Entomological Research* 99: 619-627.

MATHENGE, C.W., HOLFORD, P., HOFFMANN, J.H., ZIMMERMANN, H.G., SPOONER-HART, R. & BEATTIE, G.A.C. 2010a. Determination of biotypes of *Dactylopius tomentosus* (Hemiptera: Dactylopiidae) and insights into the taxonomic relationships of their hosts, *Cylindropuntia* spp. *Bulletin of Entomological Research* 100: 347-358.

MATHENGE, C.W., HOLFORD, P., HOFFMANN, J.H., ZIMMERMANN, H.G., SPOONER-HART, R. & BEATTIE G.A.C. 2010b. Hybridization between *Dactylopius tomentosus* (Hemiptera: Dactylopiidae) biotypes and its effects on host specificity. *Bulletin of Entomological Research* 100: 331-338.

MATTHEWS, D.L. & MAHARAJH, B.V. 2009. *Adaina primulacea* Meyrick, 1929: a gall-inducing plume moth of Siam weed from South Florida and the Neotropics. *Tropical Lepidoptera Research* 19: 64-70.

McCONNACHIE, A.J. 2015a. Host range and risk assessment of *Zygogramma bicolorata*, a defoliating agent released in South Africa for the biological control of *Parthenium hysterophorus*. *Biocontrol Science and Technology* 25: 975-991.

McCONNACHIE, A.J. 2015b. Host range tests cast doubt on the suitability of *Epiblema strenuana* as a biological control agent for *Parthenium hysterophorus* in Africa. *BioControl* 60: 1-9.

McCONNACHIE, A.J., HILL, M.P. & BYRNE, M.J. 2004. Field assessment of a frond-feeding weevil, a successful biological control agent of red water fern, *Azolla filiculoides*, in southern Africa. *Biological Control* 29: 326-331.

McCONNACHIE, A.J., RETIEF, E., HENDERSON, L. & Mc KAY, F. 2011. The initiation of a biological control programme against pompom weed, *Campuloclinium macrocephalum* (Less.) DC. (Asteraceae), in South Africa. *African Entomology* 19: 258-268.

Mc KAY, F. & GANDOLFO, D. 2007. Phytophagous insects associated with the reproductive structures of mesquite (*Prosopis* spp.) in Argentina and their potential as biocontrol agents in South Africa. *African Entomology* 15: 121-131.

Mc KAY, F., OLERIO, M., FOURIE, A. & SIMELANE, D. 2010. Natural enemies of balloon vine, *Cardiospermum grandiflorum* Swartz (Sapindaceae), in Argentina and their potential use as biological control agents in South Africa. *International Journal of Insect Science* 30: 67-76.

- MORAN, P.J. & GOOLSBY, J.A. 2009. Biology of the galling wasp *Tetramesa romana*, a biological control agent of giant reed. *Biological Control* 49: 169-179.
- MORAN, P.J. & GOOLSBY, J.A. 2010. Biology of the armored scale *Rhizaspidotus donacis* (Hemiptera: Diaspididae), a candidate agent for biological control of giant reed. *Annals of the Entomological Society of America* 103: 252-263.
- MORAN, V.C. & ANNECKE, D.P. 1979. Critical reviews of biological control in South Africa. 3. The jointed cactus, *Opuntia aurantiaca* Lindley. *Journal of the Entomological Society of southern Africa* 42: 299-329.
- MORAN, V.C. & ZIMMERMANN, H.G. 1991a. Biological control of jointed cactus, *Opuntia aurantiaca* (Cactaceae), in South Africa. *Agriculture, Ecosystems and Environment* 37: 5-27.
- MORAN, V.C. & ZIMMERMANN, H.G. 1991b. Biological control of cactus weeds of minor importance in South Africa. *Agriculture, Ecosystems and Environment* 37: 37-55.
- MORRIS, M.J. 1987. Biology of the *Acacia* gall rust, *Uromycladium tepperianum*. *Plant Pathology* 36: 100-106.
- MORRIS, M.J. 1989. Host specificity studies of a leaf spot fungus, *Phaeoramularia* sp., for the biological control of Crofton Weed (*Ageratina adenophora*) in South Africa. *Phytophylactica* 21: 281-283.
- MORRIS, M.J. 1991. The use of plant pathogens for biological weed control in South Africa. *Agriculture, Ecosystems and Environment* 37: 239-255.
- MORRIS, M.J. 1997. Impact of the gall-forming rust fungus *Uromycladium tepperianum* on the invasive tree *Acacia saligna* in South Africa. *Biological Control* 10: 75-82.
- MORRIS, M.J. 1999. The contribution of the gall-forming rust fungus *Uromycladium tepperianum* (Sacc.) McAlp. to the biological control of *Acacia saligna* (Labill.) Wendl. (Fabaceae) in South Africa. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology Memoir No. 1*. 125-128.
- MORRIS, M.J., WOOD, A., & DEN BREEYEN, A. 1999. Plant pathogens and biological control of weeds in South Africa: a review of projects and progress during the last decade. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology Memoir No. 1*. 129-137.
- MOUND, L.A. & PEYRERA, V. 2008. *Liothrips tractabilis* sp.n. (Thysanoptera: Phlaeothripinae) from Argentina. *Neotropical Entomology* 37: 63-67.
- MPHEPHU, T.E., OLCKERS T. & SIMELANE, D.O. 2017. The tortoise beetle *Physonota maculiventris* (Chrysomelidae: Cassidinae) is suitable for release against the weedy Mexican sunflower *Tithonia diversifolia* (Asteraceae) in South Africa. *Biocontrol Science and Technology* 27: 510-524.
- MUKWEVHO, L., OLCKERS, T. & SIMELANE, D.O. 2017. Establishment, dispersal and impact of the flower-galling mite *Aceria lantanae* (Acari: Trombidiformes: Eriophyidae) on *Lantana camara* (Verbenaceae) in South Africa. *Biological Control* 107: 33-40.

NZAMA, S., OLCKERS, T. & ZACHARIADES, C. 2014a. Is oviposition and larval damage by the leaf-mining fly *Calycomyza eupatorivora* (Agromyzidae) on its target weed, *Chromolaena odorata* (Asteraceae), restricted by leaf-quality preferences? *Biocontrol Science and Technology* 24: 680-689.

NZAMA, S., OLCKERS, T. & ZACHARIADES, C. 2014b. Seasonal activity, habitat preferences and larval mortality of the leaf-mining fly *Calycomyza eupatorivora* (Agromyzidae), a biological control agent established on *Chromolaena odorata* (Asteraceae) in South Africa. *Biocontrol Science and Technology* 24: 1297-1307.

OBERHOLZER, I.G., MAFOKOANE, D.L. & HILL, M.P. 2007. The biology and laboratory host range of the weevil *Listronotus marginicollis* (Hustache) (Coleoptera: Curculionidae), a natural enemy of the invasive aquatic weed *Myriophyllum aquaticum* (Velloso) Verde (Haloragaceae) (Parrot's Feather). WRC Report No. KV 180/07. Water Research Commission, Pretoria. 14 pp.

OLCKERS, T. 1995. Indigenous parasitoids inhibit the establishment of a gall-forming moth imported for the biological control of *Solanum elaeagnifolium* Cavanilles (Solanaceae) in South Africa. *African Entomology* 3: 85-87.

OLCKERS, T. 1998. Biology and host range of *Platyphora semiviridis*, a leaf beetle evaluated as a potential biological control agent for *Solanum mauritianum* in South Africa. *BioControl* 43: 225-239.

OLCKERS, T. 1999. Biological control of *Solanum mauritianum* Scopoli (Solanaceae) in South Africa: a review of candidate agents, progress and future prospects. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology* Memoir No. 1. 65-73.

OLCKERS, T. 2000a. Biology, host specificity and risk assessment of *Gargaphia decoris*, the first agent to be released in South Africa for the biological control of the invasive tree *Solanum mauritianum*. *BioControl* 45: 373-388.

OLCKERS, T. 2000b. Biology and physiological host range of four species of *Platyphora* Gistel (Coleoptera: Chrysomelidae) associated with *Solanum mauritianum* Scop. (Solanaceae) in South Africa. *The Coleopterists Bulletin* 54: 497-510.

OLCKERS, T. 2003. Assessing the risks associated with the release of a flowerbud weevil, *Anthonomus santacruzi*, against the invasive tree *Solanum mauritianum* in South Africa. *Biological Control* 28: 302-312.

OLCKERS, T. 2004. Biology, host specificity and risk assessment of the leaf-mining flea beetle, *Acallepitrax* sp. nov., a candidate agent for the biological control of the invasive tree *Solanum mauritianum* in South Africa. *BioControl* 49: 323-339.

OLCKERS, T. 2008. *Anthonomus santacruzi* Hustache (Curculionidae), a new biological control agent for bugweed, *Solanum mauritianum* Scopoli, in South Africa, poses no risks to cotton production. *African Entomology* 16: 137-139.

OLCKERS, T. 2009. 20. *Solanum mauritianum* Scopoli (Solanaceae). In: Muniappan, R., Reddy, G.V.P. & Raman, A. (Eds) *Biological Control of Tropical Weeds using Arthropods*. 408-422. Cambridge University Press, Cambridge.

OLCKERS, T. 2011a. Biological control of *Leucaena leucocephala* (Lam.) de Wit (Fabaceae) in South Africa: a tale of opportunism, seed-feeders and unanswered questions. *African Entomology* 19: 356-365.

OLCKERS, T. 2011b. Biological control of *Solanum mauritianum* Scop. (Solanaceae) in South Africa: will perseverance pay off? *African Entomology* 19: 416-426.

OLCKERS, T. & BOREA, C.K. 2009. Assessing the risks of releasing a sap-sucking lace bug, *Gargaphia decoris*, against the invasive tree *Solanum mauritianum* in New Zealand. *BioControl* 54: 143-154.

OLCKERS, T. & HULLEY, P.E. 1994. Resolving ambiguous results of host-specificity tests: the case of two *Leptinotarsa* species (Coleoptera: Chrysomelidae) for biological control of *Solanum elaeagnifolium* Cavanilles (Solanaceae) in South Africa. *African Entomology* 2: 137-144.

OLCKERS, T. & LOTTER, W.D. 2004. Possible non-target feeding by the bugweed lace bug *Gargaphia decoris* (Tingidae) in South Africa: field evaluations support predictions of laboratory host-specificity tests. *African Entomology* 12: 283-285.

OLCKERS, T. & ZIMMERMANN, H.G. 1991. Biological control of silverleaf nightshade, *Solanum elaeagnifolium*, and bugweed, *Solanum mauritianum* (Solanaceae) in South Africa. *Agriculture, Ecosystems and Environment* 37: 137-155.

OLCKERS, T., HOFFMANN, J.H., MORAN, V.C., IMPSON, F.A.C. & HILL, M.P. 1999. The initiation of biological control programmes against two herbaceous *Solanum* weed species in South Africa. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology* Memoir No. 1. 55-63.

OLCKERS, T., ZIMMERMANN, H.G. & HOFFMANN, J.H. 1995. Interpreting ambiguous results of host-specificity tests in biological control of weeds: assessment of two *Leptinotarsa* species (Chrysomelidae) for the control of *Solanum elaeagnifolium* (Solanaceae) in South Africa. *Biological Control* 5: 336-344.

PARK, I.J. 2010. Biology of *Asphondylia prosopidis* complex (Diptera: Cecidomyiidae) and fungal associates: Potential biological control candidates for South African mesquite. M.S. Thesis. New Mexico State University. 52 pp.

PATERSON, I.D., HILL, M.P. & DOWNIE, D.A. 2012. The effect of host plant intraspecific genetic variation on the fitness of a monophagous biological control agent. *Biocontrol Science and Technology*. 22(5): 513-525.

PATERSON, I.D., HOFFMANN, J.H., KLEIN, H., MATHENGE, C.W., NESER, S. & ZIMMERMANN, H.G. 2011. Biological control of Cactaceae in South Africa. *African Entomology* 19: 230-246.

PATERSON, I.D., MANGAN, R., DOWNIE, D.A., COETZEE, J.A., HILL, M.P., BURKE, A.M., DOWNEY, P.O., HENRY, T.J. & COMPTON, S. 2016. Two in one: cryptic species discovered in biological control agent populations using molecular data and interbreeding experiments. *Ecology and Evolution* 6: 6139-6150.

PATERSON, I.D., MDODANA, L.A., MPEKULA, O., MABUNDA, D.B.X. & HILL, M.P. 2014a. A promising biological control agent for the control of the invasive alien plant, *Pereskia aculeata* Miller (Cactaceae), in South Africa. *Biocontrol Science and Technology*. 24(10): 1083-1095.

- PATERSON, I.D., VITORINO, M.D, DE CRISTO S.C., MARTIN, G. & HILL, M.P. 2014b. Prioritization of potential agents for the biological control of the invasive alien weed, *Pereskia aculeata* (Cactaceae), in South Africa. *Biocontrol Science and Technology*. 24: 407-425.
- PEDROSA-MACEDO, J.H., OLCKERS, T.VITORINO, M.D. & M.G. CAXAMBU. 2003. Phytophagous arthropods associated with *Solanum mauritianum* Scopoli (Solanaceae) in the First Plateau of Paraná, Brazil: a cooperative project on biological control of weeds between Brazil and South Africa. *Neotropical Entomology* 32: 519-522.
- PHENYE, M.S. & SIMELANE, D.O. 2005. Arid-adapted Mexican leafhopper, *Barela parvisaccata* (Hemiptera: Cicadellidae), unsuitable for biocontrol of *Lantana camara* (Verbenaceae) in South Africa. *Biocontrol Science and Technology* 15: 615-621.
- RENTERIA, B., ELLISON, J.L. & ELLISON, C.A. 2004. Potential biological control of *Lantana camara* in the Galapagos using the rust *Puccinia lantanae*. *Sida* 21: 1009-1017.
- RETIEF, E. 2010a. Leaf-spot fungus fails to establish on *Lantana camara* in South Africa. *Biocontrol News and Information* 31: 21N-22N.
- RETIEF, E. 2010b. The rust fungus, *Prosopidium tuberculatum*, rejected as a possible biocontrol agent for *Lantana camara* in South Africa. *Plant Protection News* 85: 15-16.
- RETIEF, E., VAN ROOI, C. & DEN BREEYEN, A. 2016. Environmental requirements and host-specificity of *Puccinia eupatorii*, a potential biocontrol agent of *Campuloclinium macrocephalum* in South Africa. *Australasian Plant Pathology* 45 (2): 135-144.
- ROBERTS, A. 2006. Biological control of alien invasive mesquite species (*Prosopis*) in South Africa: the role of introduced seed-feeding bruchids. Unpubl. PhD thesis, University of Cape Town.
- ROBINSON, M.A., ZACHARIADES, C., ROBINSON, D.E., COHEN, J.E. & YOUNGER, N. 2013. Field host range of *Melanagromyza eupatoriella* in Jamaica: an insect with potential as a biological control agent on *Chromolaena odorata* in South Africa. In: Zachariades, C., Strathie, L.W., Day, M.D. & Muniappan, R. (eds) *Proceedings of the Eighth International Workshop on Biological Control of Chromolaena odorata and other Eupatorieae*, Nairobi, Kenya, 1-2 November 2010. ARC-PPRI, Pretoria, South Africa, pp. 102-109.
- SHARRATT, M.E. & OLCKERS, T. 2012. The biological control agent *Acanthoscelides macrophthalmus* (Chrysomelidae: Bruchinae) inflicts levels of seed damage on its target, the invasive tree *Leucaena leucocephala* (Fabaceae), in the KwaZulu-Natal coastal region of South Africa. *African Entomology* 20(1): 44-51.
- SHOBA, Z. & OLCKERS, T. 2010. Re-assessment of the biology and host range of *Acanthoscelides macrophthalmus* (Chrysomelidae: Bruchinae), a seed-feeding beetle released for the biological control of *Leucaena leucocephala* in South Africa. *African Entomology* 18 Supplement No. 1: 1-9.
- SIMELANE, D.O. 2002. Biology and host range of *Ophiomyia camarae*, a biological control agent for *Lantana camara* in South Africa. *BioControl* 47: 575-585.
- SIMELANE, D.O. 2005. Biological control of *Lantana camara* in South Africa: targeting a different niche with a root-feeding agent, *Longitarsus* sp. *BioControl* 50: 375-387.

- SIMELANE, D.O. 2010. Potential impact of an introduced root-feeding flea beetle, *Longitarsus bethae*, on the growth and reproduction of an invasive weed, *Lantana camara*. *Biological Control* 54: 114-118.
- SIMELANE, D.O., FOURIE, A. & MAWELA, K.V. 2011a. Prospective agents for the biological control of *Cardiospermum grandiflorum* Sw. (Sapindaceae) in South Africa. *African Entomology* 19: 269-277.
- SIMELANE, D.O., MAWELA, K.V. & FOURIE, A. 2011b. Prospective agents for the biological control of *Tithonia rotundifolia* (Mill.) S.F.Blake and *Tithonia diversifolia* (Hemsl.) A.Gray (Asteraceae), in South Africa. *African Entomology* 19: 443-450.
- SIMELANE, D.O. & PHENYE, M.S. 2005. Suppression of growth and reproductive capacity of the weed *Lantana camara* (Verbenaceae) by *Ophiomyia camarae* (Diptera: Agromyzidae) and *Teleonemia scrupulosa* (Heteroptera: Tingidae). *Biocontrol Science and Technology* 15: 153-163.
- SINGH, D. AND OLCKERS, T. 2017. Climate constrains the establishment and proliferation of *Anthonomus santacruzi*, a flower-feeding biological control agent of the invasive weed *Solanum mauritianum* in South Africa. *Biocontrol Science and Technology* 27: 475-484.
- SOLIS, M.A., METZ, M. & ZACHARIADES, C. 2008. Identity and generic placement of *Phestinia costella* Hampson (Lepidoptera: Pyralidae: Phycitinae) reared on the invasive plant *Chromolaena odorata* (L.) R.M. King & H. Rob. (Asteraceae). *Proceedings of the Entomological Society of Washington* 110: 292-301.
- SPARKS, H.E. 1999. The initiation of a biological control programme against *Macfadyena unguis-cati* (L.) Gentry (Bignoniaceae) in South Africa. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology* Memoir No. 1. 153-157.
- STRATHIE, L.W. 1995. Ovipositional behaviour and efficacy of *Neltumius arizonensis*, a potential biological control agent of *Prosopis* weeds in South Africa. MSc thesis, University of Cape Town.
- STRATHIE, L.W., McCONNACHIE, A.J. & RETIEF, E. 2011. Initiation of biological control against *Parthenium hysterophorus* L. (Asteraceae) in South Africa. *African Entomology* 19: 378-392.
- STRATHIE, L. W. & ZACHARIADES, C. 2002. Biological control of *Chromolaena odorata* in South Africa: developments in research and implementation. In: Zachariades, C., Muniappan, R. & Strathie, L.W. (Eds) *Proceedings of the Fifth International Workshop on Biological Control and Management of Chromolaena odorata, Pretoria, South Africa*. 74-79. ARC-PPRI, Pretoria.
- STRATHIE, L.W. & ZACHARIADES, C. 2004. Insects for the biological control of *Chromolaena odorata*: surveys in the northern Caribbean and efforts undertaken in South Africa. In: Day, M.D. & McFadyen, R.E.C. (Eds) *Proceedings of the Sixth International Workshop on Biological Control and Management of Chromolaena*. May 2003. *ACIAR Technical Reports* 55: 45-52.
- STRATHIE, L.W., ZACHARIADES, C., GAREEB, M. & SHARP, D. 2007. Establishment of two biocontrol agents on *Chromolaena odorata* in South Africa. In: Lai, Po-Yung, Reddy, G.V.P. & Muniappan, R. (Eds) *Proceedings of the Seventh International Workshop on Biological Control and Management of Chromolaena odorata and Mikania micrantha, Taiwan, September 12 to 15, 2006*. 81-93. National Pingtung University of Science and Technology (NPUST), Taiwan.

- THOMAS, S.E., ELLISON, C.A. & TOMLEY, A.J. 2006. Studies on the rust *Prospodium tuberculatum*, a new classical biological control agent released against the invasive alien weed *Lantana camara* in Australia. *Australasian Plant Pathology* 35: 321-328.
- TRACY, J.L. & ROBBINS, T.O. 2009. Taxonomic revision and biogeography of the *Tamarix*-feeding *Diorhabda elongata* (Brullé, 1832) species group (Coleoptera: Chrysomelidae: Galerucinae: Galerucini) and analysis of their potential in biological control of tamarisk. *Zootaxa* 2101: 1-152.
- TRUJILLO, E.E. 1995. *Septoria* leaf spot of lantana from Ecuador: a potential biological control for bush lantana in forests of Hawaii. *Plant Disease* 79: 819-821.
- URBAN, A.J., MPEDI, P.F., NESER, S. & CRAEMER, C. 2001. Potential of the flower gall mite, *Aceria lantanae* (Cook) (Acari: Eriophyidae), for biocontrol of the noxious weed, *Lantana camara* L. (Verbenaceae). In: Olckers, T. & Brothers, D.J. (Eds) *Proceedings of the 13th Entomological Congress of the Entomological Society of Southern Africa*, Pietermaritzburg, 2-4 July 2001. 67-68.
- URBAN, A.J., NESER, S., & MPEDI, P. 2011a. Lantana flower gall mite: established, spreading and making an impact. *Plant Protection News* 86: 1-2.
- URBAN, A.J., SIMELANE, D.O., RETIEF, E., HEYSTEK, F., WILLIAMS, H.E. & MADIRE, L.G. 2011b. The invasive '*Lantana camara* L.' hybrid complex (Verbenaceae): a review of research into its identity and biological control in South Africa. *African Entomology* 19: 315-348.
- UYI, O.O., HILL, M.P. & ZACHARIADES, C. 2014b. Variation in host plant has no effect on the performance and fitness-related traits of the specialist herbivore *Pareuchaetes insulata*. *Entomologia Experimentalis et Applicata* 153: 64-75.
- UYI, O.O., ZACHARIADES, C. & HILL, M.P. 2014a. The life history traits of the arctiine moth *Pareuchaetes insulata*, a biological control agent of *Chromolaena odorata* in South Africa. *African Entomology* 22: 611-624.
- UYI, O.O., ZACHARIADES, C., HILL, M.P. & CONLONG, D. 2015. The nocturnal larvae of a specialist folivore perform better on *Chromolaena odorata* leaves from a shaded environment. *Entomologia Experimentalis et Applicata* 156: 187-199.
- UYI, O.O., ZACHARIADES, C. & HILL, M.P. 2016a. Nitrogen fertilisation improves growth of *Chromolaena odorata* (Asteraceae) and the performance of the biological control agent, *Pareuchaetes insulata* (Erebidae). *Biocontrol Science and Technology* 26: 373-385.
- UYI, O.O., ZACHARIADES, C., HILL, M.P. & McCONNACHIE, A.J. 2016b. Temperature-dependent performance and potential distribution of *Pareuchaetes insulata*, a biological control agent of *Chromolaena odorata* in South Africa. *BioControl* 61: 815-825.
- UYI, O.O., ZACHARIADES, C., MARAIS, E., HILL, M.P. 2017. Reduced mobility but high survival: thermal tolerance and locomotor response of the specialist herbivore, *Pareuchaetes insulata* (Walker) (Lepidoptera: Erebidae), to low temperatures. *Bulletin of Entomological Research* 107: 448-457.
- VAN DER WESTHUIZEN, L. 2006. The evaluation of *Phenrica* sp.2 (Coleoptera: Chrysomelidae: Alticinae), as a possible biological control agent for *Anredera cordifolia* (Ten.) Steenis in South Africa. MSc thesis, Rhodes University, Grahamstown.

- VAN DER WESTHUIZEN, L. 2011. Initiation of a biological control programme against Madeira vine, *Anredera cordifolia* (Ten.) Steenis (Basellaceae), in South Africa. *African Entomology* 19: 217-222.
- VAN DER WESTHUIZEN, L. & MPEDI, P. 2011. The initiation of a biological control programme against *Argemone mexicana* L. and *Argemone ochroleuca* Sweet subsp. *ochroleuca* (Papaveraceae) in South Africa. *African Entomology* 19: 223-229.
- VAN KLINKEN, R.D., HOFFMANN, J.H., ZIMMERMANN, H.G. & ROBERTS, A.P. 2009. *Prosopis* species (Leguminosae). In: Muniappan, R., Reddy, G.V.P. & Raman, A. (Eds) *Biological Control of Tropical Weeds using Arthropods*. 353-377. Cambridge University Press, Cambridge.
- WHEELER, G.S. & CENTER, T.D. 1996. The influence of hydrilla leaf quality on larval growth and development of the biological control agent, *Hydrellia pakistanae* (Diptera: Ephydriidae). *Biological Control* 7: 1-9.
- WHEELER, G.S. & CENTER, T.D. 2001. Impact of the biological control agent *Hydrellia pakistanae* (Diptera: Ephydriidae) on the submersed aquatic weed *Hydrilla verticillata* (Hydrocharitaceae). *Biological Control* 21: 168-181.
- WILLIAMS, H.E. 2002. Life history and laboratory host range of *Charidotis auroguttata* (Boheman) (Coleoptera: Chrysomelidae), the first natural enemy released against *Macfadyena unguis-cati* (L.) Gentry (Bignoniaceae) in South Africa. *The Coleopterists Bulletin* 56: 299-307.
- WILLIAMS, H.E. 2004. Unsuitability of *Charidotis pygmaea* for biocontrol of *Lantana camara* in Africa. *BioControl* 49: 211-223.
- WILLIAMS, H.E. 2005. The potential impact of the trimorphic flea-beetle, *Alagoasa extrema*, on a selected weedy variety of *Lantana camara*. *BioControl* 50: 647-655.
- WILLIAMS, H.E. 2006. Preference and performance of the polymorphic flea-beetle, *Alagoasa extrema*, on selected, weedy *Lantana camara* varieties. *BioControl* 51: 127-138.
- WILLIAMS, H.E. & MADIRE, L.G. 2008. Biology, host range and varietal preference of the leaf-feeding geometrid, *Leptostales ignifera*, a potential biocontrol agent for *Lantana camara* in South Africa, under laboratory conditions. *BioControl* 53: 957-969.
- WILLIAMS, H.E., NESER, S. & MADIRE, L.G. 2008. Candidates for biocontrol of *Macfadyena unguis-cati* in South Africa: biology, host ranges and potential impact of *Carvalhotingis visenda* and *Carvalhotingis hollandi* under quarantine conditions. *BioControl* 53: 945-956.
- WOOD, A.R. 1999. *The potential biological control agents of Cladophora glomerata that occur in irrigation schemes in South Africa*. WRC Report No 669/1/99. Water Research Commission, Pretoria. 34 pp.
- WOOD, A.R. 2003. *The potential of aquatic Pythium species for the biological control of Cladophora glomerata in irrigation schemes in South Africa*. WRC Report No. 918/1/03. Water Research Commission, Pretoria. 39 pp.
- WOOD, A.R. 2008. Host-specificity testing of *Prospodium transformans* (Uredinales: Uropyxidaceae), a biological control agent for use against *Tecoma stans* var. *stans* (Bignoniaceae). In: Julien, M.H.,

Sforza, R., Bon, M.C., Evans, H.C., Hatcher, P.E., Hinz, H.L. & Rector, B.G. (Eds.) *Proceedings of the XII International Symposium on Biological Control of Weeds*: 345-348. CAB International, Wallingford, UK.

WOOD, A.R. & GINNS, J. 2006. A new dieback disease of *Acacia cyclops* in South Africa caused by *Pseudolagarobasidium acaciicola* sp.nov. *Canadian Journal of Botany* 84: 750-758.

WOOD, A.R. & MORRIS, M.J. 2007. Impact of the gall-forming rust fungus *Uromycladium tepperianum* on the invasive tree *Acacia saligna* in South Africa: 15 years of monitoring. *Biological Control* 41: 68-77.

WOOD, A.R. & SCHOLLER, M. 2002. *Puccinia abrupta* var. *parthenicola* on *Parthenium hysterophorus* in Southern Africa. *Plant Disease* 86: 327.

ZACHARIADES, C., HOFFMANN, J.H. & ROBERTS, A. 2011a. Biological control of mesquite (*Prosopis* species) (Fabaceae) in South Africa. *African Entomology* 19: 402-415.

ZACHARIADES, C., STRATHIE, L., DELGADO, O. & RETIEF, E. 2007. Pre-release research on biocontrol agents for chromolaena in South Africa. In: Lai, Po-Yung, Reddy, G.V.P. & Muniappan, R. (Eds) *Proceedings of the Seventh International Workshop on Biological Control and Management of Chromolaena odorata and Mikania micrantha, Taiwan, September 12 to 15, 2006*. 68-80. National Pingtung University of Science and Technology (NPUST), Taiwan.

ZACHARIADES, C., STRATHIE, L.W., RETIEF, E. & DUBE, N. 2011b. Progress towards the biological control of *Chromolaena odorata* (L.) R.M. King & H. Rob. (Asteraceae) in South Africa. In: Moran, V.C., Hoffmann, J.H. & Hill, M.P. (Eds) *Biological Control of Invasive Alien Plants in South Africa (1999–2010)*. *African Entomology* 19: 282–302.

ZACHARIADES, C., STRATHIE-KORRÛBEL, L.W., & KLUGE, R.L. 1999. The South African programme on the biological control of *Chromolaena odorata* (L.) King & Robinson (Asteraceae) using insects. In: Olckers, T. & Hill, M. P. (Eds) *Biological Control of Weeds in South Africa (1990-1999)*. *African Entomology Memoir No. 1*. 89-102.

ZACHARIADES, C., STRATHIE, L.W. & KLUGE, R.L. 2002. Biology, host-specificity and effectiveness of insects for the biocontrol of *Chromolaena odorata* in South Africa. In: Zachariades, C., Muniappan, R & Strathie, L.W. (Eds) *Proceedings of the Fifth International Workshop on Biological Control and Management of Chromolaena odorata*. 160-166. ARC-PPRI, Pretoria.

ZACHARIADES, C., UYI, O.O., DUBE, N., STRATHIE, L.W., MUIR, D., CONLONG, D.E. & ASSEFA, Y. 2016. Biological control of *Chromolaena odorata*: *Pareuchaetes insulata* spreads its wings. *Proceedings of the South African Sugar Technologists' Association* 89: 291-306.

ZIMMERMANN, H.G. 1991. Biological control of spear thistle, *Cirsium vulgare*, in South Africa. *Agriculture, Ecosystems and Environment* 37: 199-205.

ZIMMERMANN, H.G. & MORAN, V.C. 1991. Biological control of prickly pear, *Opuntia ficus-indica* (Cactaceae), in South Africa. *Agriculture, Ecosystems and Environment* 37: 29-36.

ZIMMERMANN, H., BLOEM, S. & KLEIN, H. 2004. *Biology, History, Threat, Surveillance and Control of the Cactus Moth, Cactoblastis cactorum*. Joint FAO/IAEA Programme of Nuclear Techniques in Food and Agriculture. IAEA, Vienna. 40 pp.

ZIMMERMANN, H.G., MORAN, V.C. & HOFFMANN, J.H. 2009. Invasive cactus species (Cactaceae). In: Muniappan, R., Reddy, G.V.P. & Raman, A. (Eds.) Biological Control of Tropical Weeds using Arthropods. 108-129. Cambridge University Press, Cambridge.