

Target weed species in South Africa, their region of origin, and degree of biological control achieved, where applicable (see below for definitions): natural enemies studied, their feeding guilds or association with the host plant, year of release, establishment and degree of agent damage to the weed (if established) (see below for definitions)

Target weed, region of origin, and degree of control	Natural enemy	Feeding guild	Agent status	Damage inflicted
ARACEAE				
<i>Pistia stratiotes</i> L. (water lettuce) South America Complete	<i>Neohydronomus affinis</i> Hustache (Coleoptera: Curculionidae)	Leaf and stem borer	Released 1985, established	Extensive
ASTERACEAE				
<i>Ageratina adenophora</i> (Spreng.) R.M.King & H.Rob. (formerly in <i>Eupatorium</i>) (Crofton weed) Central America Negligible	<i>Passalora ageratiniae</i> Crous & A.R. Wood (="Phaeoramularia" sp.) (anamorphic fungus; Mycosphaerellales: Mycosphaerellaceae)	Leaf spot pathogen	Released 1987, established	Moderate
	<i>Procecidochares utilis</i> Stone (Diptera: Tephritidae)	Stem galler	Released 1984, established	Moderate
<i>Ageratina riparia</i> (Regel) R.M.King & H.Rob. (formerly in <i>Eupatorium</i>) (mistflower) Central America Complete	<i>Entyloma ageratiniae</i> R.W. Barreto & H.C. Evans (Entylomatales: Entylomataceae)	Leaf pathogen	Released 1989, established	Considerable
<i>Campuloclinium macrocephalum</i> (Less.) DC. [formerly in <i>Eupatorium</i>] (pompom weed) South America Not determined	<i>Puccinia eupatorii</i> Dietel (Pucciniales: Pucciniaceae)	Leaf rust pathogen	Not released, occurs locally; also under investigation	Unknown
<i>Chromolaena odorata</i> (L.) R.M.King & H.Rob. [formerly in <i>Eupatorium</i>] (triffid weed) North, Central & South America Not determined	<i>Calycomyza eupatorivora</i> Spencer (Diptera: Agromyzidae)	Leaf miner	Released 2003, established	Moderate
	<i>Lixus aemulus</i> Petri (Coleoptera: Curculionidae)	Stem borer	Released 2011, establishment unconfirmed	
	<i>Pareuchaetes aurata</i> (Butler) (Lepidoptera: Arctiidae)	Leaf feeder	Released 1990, not established	
	<i>Pareuchaetes insulata</i> (Walker) (Lepidoptera: Arctiidae)	Leaf feeder	Released 2001, established	Considerable
	<i>Pareuchaetes pseudoinsulata</i> Rego Barros (Lepidoptera: Arctiidae)	Leaf feeder	Released 1989 & 1998, not established	
<i>Cirsium vulgare</i> (Savi) Ten. (spear thistle) Europe Negligible	<i>Rhinocyllus conicus</i> (Froelich) (Coleoptera: Curculionidae)	Seed feeder	Released 1984-1986, established	Moderate
	<i>Urophora stylata</i> (Fabricius) (Diptera: Tephritidae)	Seed feeder	Released 1984-1990, not established	
<i>Parthenium hysterophorus</i> L. (parthenium) Gulf of Mexico & South America Negligible	<i>Puccinia abrupta</i> Dietel & Holw. var. <i>partheniicola</i> (H.S. Jacks.) Parmelee (Pucciniales: Pucciniaceae)	Leaf rust pathogen	Not released, occurs locally	Unknown
	<i>Puccinia xanthii</i> Schwein. var. <i>parthenii-hysterophorae</i> Seier, H.C. Evans & Á. Romero (= <i>P. melampodii</i> ; Pucciniales: Pucciniaceae)	Leaf rust pathogen	Released 2010, establishment unconfirmed	Unknown
<i>Silybum marianum</i> (L.) Gaertn. (milk thistle) Mediterranean, Asia	<i>Rhinocyllus conicus</i> (Froelich) (Coleoptera: Curculionidae)	Seed feeder	Alternative host, released 1985, not established	
AZOLLACEAE				

<i>Azolla filiculoides</i> Lam. (red water fern) South America Complete	<i>Stenopelmus rufinasus</i> Gyllenhal (Coleoptera: Curculionidae)	Frond feeder	Released 1997, established	Extensive
BIGNONIACEAE				
<i>Macfadyena unguis-cati</i> (L.) A.H.Gentry (cat's claw creeper) Central & South America Negligible	<i>Carvalhotingis hollandi</i> Drake (Hemiptera: Tingidae)	Leaf sucker	Released 2007, established	Unknown
	<i>Carvalhotingis visenda</i> Drake & Hambleton (Hemiptera: Tingidae)	Leaf sucker	Released 2007, established	Moderate
	<i>Charidotis auroguttata</i> Boheman (Coleoptera: Chrysomelidae: Cassidinae)	Leaf feeder	Released 1999, established	Trivial
	<i>Hylaeogena (Hedwigella) jureceki</i> Obenberger (Coleoptera: Buprestidae)	Leaf miner	Released 2007, established	Unknown
	<i>Hypocosmia pyrochroma</i> Jones (Lepidoptera: Pyralidae)	Leaf feeder	Released 2010, establishment unconfirmed	Unknown
<i>Tecoma stans</i> (L.) Juss ex Kunth var. <i>stans</i> (yellow bells) North & Central America Not determined	<i>Prospodium transformans</i> (Ellis & Everh.) Cummins (Uropyxidaceae)	Leaf rust pathogen	Released 2010, establishment unconfirmed	Unknown
CACTACEAE				
<i>Austrocylindropuntia subulata</i> (Muehlenpf.) Backeb. (= <i>Opuntia exaltata</i> (Berger)) South America Not determined	<i>Cactoblastis cactorum</i> (Berg) (Lepidoptera: Pyralidae)	Cladode borer	Alternative host, established	Unknown
	<i>Hypogeococcus pungens</i> Granara de Willink (Hemiptera: Pseudococcidae)	Stem sucker	Alternative host, established	Extensive
	<i>Nealcidion cereicola</i> (Fisher) [formerly in <i>Alcidion</i>] (Coleoptera: Cerambycidae)	Stem borer	Alternative host, released 1990, established	Considerable
<i>Cylindropuntia fulgida</i> (Engelm.) F.M.Knuth var. <i>fulgida</i> (chain-fruit cholla) North & Central America Complete	<i>Dactylopius tomentosus</i> (Lamark), 'cholla' biotype (Hemiptera: Dactylopiidae)	Cladode sucker	Released 2008, established	Extensive
	<i>Dactylopius tomentosus</i> (Lamark), 'imbricata' biotype (Hemiptera: Dactylopiidae)	Cladode sucker	Alternative host, released 1970, established	Trivial
<i>Cylindropuntia imbricata</i> (Haw.) F.M.Knuth (imbricate prickly pear) North & Central America Substantial	<i>Dactylopius tomentosus</i> (Lamark), 'imbricata' biotype (Hemiptera: Dactylopiidae)	Cladode sucker	Released 1970, established	Considerable
	<i>Metamasius spinolae</i> (Gyllenhal) (Coleoptera: Curculionidae)	Stem borer	Alternative host, released 1974, not established	
<i>Cylindropuntia leptocaulis</i> (DC.) F.M.Knuth (pencil cactus) North & Central America Complete	<i>Dactylopius tomentosus</i> (Lamark), 'imbricata' biotype, (Hemiptera: Dactylopiidae)	Cladode sucker	Alternative host, released 1977, established	Extensive
<i>Harrisia bonplandii</i> (Pfeiff.) Britton & Rose (= <i>H. balansae</i> (K.Schum.) N.P.Taylor & Zappi) South America Substantial	<i>Hypogeococcus pungens</i> Granara de Willink (Hemiptera: Pseudococcidae)	Stem sucker	Alternative host, released 2006, established	Extensive
<i>Harrisia martinii</i> (Labour.) Britton & Rose (harrisia/moon cactus)	<i>Hypogeococcus pungens</i> Granara de Willink (Hemiptera: Pseudococcidae)	Stem sucker	Released 1983, established	Extensive

South America Complete	<i>Nealcidion cereicola</i> (Fisher) [formerly in <i>Alcidion</i>] (Coleoptera: Cerambycidae)	Stem borer	Released 1990, established	Considerable
South America Substantial	<i>Cactoblastis cactorum</i> (Berg) (Lepidoptera: Pyralidae)	Cladode borer	Alternative host, established	Moderate
	<i>Dactylopius austrinus</i> De Lotto (Hemiptera: Dactylopiidae)	Cladode sucker	Released 1935, established	Extensive
	<i>Mimorista pulchellalis</i> Dyar (Lepidoptera: Crambidae)	Cladode borer	Released 1979, not established	
	<i>Nanaia</i> sp. (Lepidoptera: Pyralidae)	Cladode borer	Released 1983, not established	
	<i>Zophodia tapiacola</i> (Dyar) [formerly in <i>Tucumania</i>] (Lepidoptera: Pyralidae)	Cladode borer	Released 1976 & 1982, not established	
<i>Opuntia engelmannii</i> Salm-Dyck ex Engelm. (= <i>O. lindheimeri</i> , <i>O. tardospina</i>) (small round-leaved prickly pear) North & Central America Substantial	<i>Cactoblastis cactorum</i> (Berg) (Lepidoptera: Pyralidae)	Cladode borer	Alternative host, released 1938, established	Extensive
	<i>Dactylopius opuntiae</i> (Cockerell), 'ficus' biotype (Hemiptera: Dactylopiidae)	Cladode sucker	Alternative host, released 1938, established	Considerable
<i>Opuntia ficus-indica</i> (L.) Mill. (mission prickly pear) Central America Substantial	<i>Cactoblastis cactorum</i> (Berg) (Lepidoptera: Pyralidae)	Cladode borer	Released 1933, established	Extensive
	<i>Dactylopius opuntiae</i> (Cockerell), 'ficus' biotype (Hemiptera: Dactylopiidae)	Cladode sucker	Released 1938, established	Extensive
	<i>Lagocheirus funestus</i> (Thompson) [formerly in <i>Archagocheirus</i>] (Coleoptera: Cerambycidae)	Stem borer	Released 1943, established	Trivial
	<i>Metamasius spinolae</i> (Gyllenhal) (Coleoptera: Curculionidae)	Stem borer	Released 1948, established	Extensive
<i>Opuntia monacantha</i> Haw. (smooth/ drooping prickly pear) South America Substantial	<i>Cactoblastis cactorum</i> (Berg) (Lepidoptera: Pyralidae)	Cladode borer	Alternative host, established	Considerable
	<i>Dactylopius ceylonicus</i> (Green) (Hemiptera: Dactylopiidae)	Cladode sucker	Released 1913, established	Extensive
<i>Opuntia salmiana</i> J.Parm. ex Pfeiff. South America Substantial	<i>Cactoblastis cactorum</i> (Berg) (Lepidoptera: Pyralidae)	Cladode borer	Alternative host, established	Considerable
<i>Opuntia spinulifera</i> Salm-Dyck (large round-leaved prickly pear) Central America Not determined	<i>Cactoblastis cactorum</i> (Berg) (Lepidoptera: Pyralidae)	Cladode borer	Alternative host, established	Unknown
<i>Opuntia stricta</i> (Haw.) Haw. (Australian pest pear) North America & Caribbean Substantial	<i>Cactoblastis cactorum</i> (Berg) (Lepidoptera: Pyralidae)	Cladode borer	Alternative host, released 1980s, established	Extensive
	<i>Dactylopius opuntiae</i> (Cockerell), Australian ('stricta') biotype (Hemiptera: Dactylopiidae)	Cladode sucker	Released 1997, established	Extensive
<i>Pereskia aculeata</i> Mill. (Barbados gooseberry/ pereskia) South America & Caribbean Negligible	<i>Phenrica guerini</i> Bechyné (Coleoptera: Chrysomelidae)	Leaf feeder	Released 1991 & 1995, established	Trivial
CLUSIACEAE				
Europe & Asia Complete	<i>Hypericum perforatum</i> L. (St John's wort)	<i>Agrilus (Spiragrilus) hyperici</i> (Creutzer) (Coleoptera: Buprestidae)	Root borer	Released 1974- 1979, not established
		<i>Aplocera efformata</i> (Guenée) (Lepidoptera: Geometridae)	Leaf feeder	Released 1983, not established

	<i>Aphis chloris</i> Koch (Hemiptera: Aphididae)	Sap sucker	Released 1982, not established	
	<i>Chrysolina hyperici</i> (Forster) (Coleoptera: Chrysomelidae: Chrysomelinae)	Leaf feeder	Released 1960-1973, not established	
	<i>Chrysolina quadrigemina</i> Suffrian (Coleoptera: Chrysomelidae: Chrysomelinae)	Leaf feeder	Released 1960-1973, established	Extensive
	<i>Zeuxidiplosis giardi</i> (Kieffer) (Diptera: Cecidomyiidae)	Shoot-tip galler	Released 1972, established	Moderate
FABACEAE				
<i>Caesalpinia decapetala</i> (Roth) Alston (Mauritius thorn) Asia Negligible	<i>Sulcobruchus subsuturalis</i> (Pic) (= <i>S. bakeri</i> Kingsolver) (Coleoptera: Chrysomelidae: Bruchinae)	Seed feeder	Released 1999, established	Trivial
<i>Gleditsia triacanthos</i> L. (honey locust) North America Not determined	<i>Megabruchidius tonkineus</i> (Pic) (Coleoptera: Chrysomelidae: Bruchinae)	Seed feeder	Not released, occurs locally	Unknown
<i>Leucaena leucocephala</i> (Lam.) de Wit (leucaena) Tropical America Negligible	<i>Acanthoscelides macrophthalmus</i> (Schaeffer) (Coleoptera: Chrysomelidae: Bruchinae)	Seed feeder	Released 1999, established	Trivial
<i>Prosopis</i> L. species (mesquite/ prosopis) North & Central America Negligible	<i>Algarobius bottimeri</i> Kingsolver (Coleoptera: Chrysomelidae: Bruchinae)	Seed feeder	Released 1990, not established	
	<i>Algarobius prosopis</i> (LeConte) (Coleoptera: Chrysomelidae: Bruchinae)	Seed feeder	Released 1987, established	Considerable
	<i>Neltumius arizonensis</i> (Schaeffer) (Coleoptera: Chrysomelidae: Bruchinae)	Seed feeder	Released 1993, established	Unknown
<i>Sesbania punicea</i> (Cav.) Benth. (red sesbania) South America Complete	<i>Neodiplogrammus quadrivittatus</i> (Olivier) (Coleoptera: Curculionidae)	Stem borer	Released 1984, established	Extensive
	<i>Rhyssomatus marginatus</i> Fähraeus (Coleoptera: Curculionidae)	Seed feeder	Released 1984, established	Extensive
	<i>Trichapion lativentre</i> (Béguin-Billecocq) (Coleoptera: Brentidae: Apioninae)	Flowerbud feeder	Released 1970s, established, present before release	Extensive
HALORAGACEAE				
<i>Myriophyllum aquaticum</i> (Vell.) Verdc. (parrot's feather) South America Substantial	<i>Lysathia</i> sp. (Coleoptera: Chrysomelidae)	Leaf feeder	Released 1994, established	Extensive
	<i>Xanthomonas campestris</i> (Pammel) Dawson (?) (Proteobacteria: Xanthomonadales; Xanthomonadaceae)	Bacterial wilt	Not released, occurs locally; rejected	
MIMOSACEAE				
<i>Acacia baileyana</i> F.Muell. (Bailey's wattle) Australia Negligible	<i>Melanterius maculatus</i> Lea (Coleoptera: Curculionidae)	Seed feeder	Alternative host, released 2006 & 2009, established	Moderate
<i>Acacia cyclops</i> A.Cunn. ex G.Don (red eye/ rooikrans) Australia Substantial	<i>Calonectria scoparia</i> Ribeiro & Matsuoka ex Peerally (Hypocreales: Nectriaceae) (anamorphic fungus: <i>Cylindrocladium candelabrum</i>)	Leaf spot pathogen	Not released, occurs locally; rejected	
	<i>Dasineura dielsi</i> Rübsaamen (Diptera: Cecidomyiidae)	Flower galler	Released 2002, established	Extensive
	<i>Melanterius servulus</i> Pascoe (Coleoptera: Curculionidae)	Seed feeder	Alternative host, released 1991 &	Considerable

			1994, established	
	<i>Pseudolagarobasidium acaciicola</i> Ginns (Polyporales: Phanerochaetaceae)	Die-back pathogen	Not released, occurs locally; under investigation	Unknown
<i>Acacia dealbata</i> Link (silver wattle) Australia Not determined	<i>Melanterius maculatus</i> Lea (Coleoptera: Curculionidae)	Seed feeder	Alternative host, released 1994- 2007, established	Moderate
<i>Acacia decurrens</i> (Wendl.) Willd. (green wattle) Australia Negligible	<i>Melanterius maculatus</i> Lea (Coleoptera: Curculionidae)	Seed feeder	Alternative host, released 2001- 2007, established	Moderate
<i>Acacia longifolia</i> (Andr.) Willd. (long-leaved wattle) Australia Substantial	<i>Melanterius ventralis</i> Lea (Coleoptera: Curculionidae)	Seed feeder	Released 1985, established	Extensive
	<i>Trichilogaster acaciaelongifoliae</i> (Froggatt) (Hymenoptera: Pteromalidae)	Bud galler	Released 1982, established	Extensive
<i>Acacia mearnsii</i> De Wild. (black wattle) Australia Not determined	<i>Ceratocystis albofundus</i> M.J.Wingf., De Beer & M.J.Morris (Microascales: Ceratocystidaceae)	Gummosis pathogen	Not released, occurs locally; shelved	
	<i>Cylindrobasidium laeve</i> (Pers.) Chamuris (Polyporales: Physalacriaceae)	Early succession wood rot pathogen, prevents coppice growth	Not released, occurs locally; developed as cut stump inoculant "Stumpout"	
	<i>Dasineura rubiformis</i> Kolesik (Diptera: Cecidomyiidae)	Flower galler	Present before release, established	Extensive (but still localized)
	<i>Melanterius maculatus</i> Lea (Coleoptera: Curculionidae)	Seed feeder	Released 1993- 1996, established	Moderate
<i>Acacia melanoxylon</i> R.Br. (Australian blackwood) Australia Substantial	<i>Melanterius acaciae</i> Lea (Coleoptera: Curculionidae)	Seed feeder	Released 1986, established	Extensive
<i>Acacia podalyriifolia</i> A.Cunn. ex G.Don (pearl acacia) Australia Not determined	<i>Melanterius maculatus</i> Lea (Coleoptera: Curculionidae)	Seed feeder	Alternative host, released 2008, established	Unknown
<i>Acacia pycnantha</i> Benth. (golden wattle) Australia Substantial	<i>Cylindrobasidium laeve</i> (Pers.) Chamuris (Polyporales: Physalacriaceae)	Early succession wood rot fungus	Alternative host; not released, occurs locally; developed as cut stump inoculant "Stumpout", but never used	
	<i>Melanterius maculatus</i> Lea (Coleoptera: Curculionidae)	Seed feeder	Alternative host, released 2005- 2007, established	Moderate
	<i>Trichilogaster signiventris</i> (Girault) (Hymenoptera: Pteromalidae)	Bud galler	Released 1987 & 1992, established	Considerable / Extensive
<i>Acacia saligna</i> (Labill.) H.L.Wendl. (Port Jackson willow) Australia Substantial	<i>Melanterius compactus</i> Lea (Coleoptera: Curculionidae)	Seed feeder	Released 2001 & 2003, established	Considerable
	<i>Uromycladium tepperianum</i> (Sacc.) McAlpine (Pucciniales: Pileolariaceae)	Gall former	Released 1987, established	Extensive
<i>Paraserianthes lophantha</i> (Willd.) Nielsen [formerly in <i>Albizia</i>] (stink bean) Australia	<i>Melanterius servulus</i> Pascoe (Coleoptera: Curculionidae)	Seed feeder	Released 1989, established	Considerable

Substantial				
MYRTACEAE				
<i>Leptospermum laevigatum</i> (Gaertn.) F.Muell. (Australian myrtle) Australia Negligible	<i>Aristaea thalassias</i> (Meyrick) [formerly in <i>Parectopa</i>] (Lepidoptera: Gracillariidae)	Leaf feeder	Released 1996, established	Considerable
	<i>Dasineura strobila</i> Dorchin (Diptera: Cecidomyiidae)	Bud galler	Released 1994, established, present before release	Considerable
PONTEDERIACEAE				
<i>Eichhornia crassipes</i> (C.Mart.) Solms (water hyacinth) South America Substantial	<i>Acremonium zonatum</i> (Sawada) W.Gams (anamorphic fungus; Hypocreales)	Leaf pathogen	Not released, occurs locally	
	<i>Alternaria eichhorniae</i> Nag-Raj & Ponnappa (anamorphic fungus; Pleosporales: Pleosporaceae)	Leaf pathogen	Not released, occurs locally; shelved	
	<i>Cercospora piaropi</i> Tharp. (anamorphic fungus; Mycosphaerellales: Mycosphaerellaceae)	Leaf pathogen	Not released, occurs locally	
	<i>Cercospora rodmanii</i> Conway (anamorphic fungus; Mycosphaerellales: Mycosphaerellaceae)	Leaf pathogen	Released 1992, established	Considerable
	<i>Cornops aquaticum</i> (Brüner) (Orthoptera: Acrididae: Leptysminae)	Leaf feeder	Released 2011, establishment unconfirmed	
	<i>Eccritotarsus catarinensis</i> (Carvalho) (Hemiptera: Miridae)	Leaf sucker	Released 1996, established	Considerable
	<i>Neochetina bruchi</i> Hustache (Coleoptera: Curculionidae)	Stem borer	Released 1990 & 1996, established	Considerable
	<i>Neochetina eichhorniae</i> Warner (Coleoptera: Curculionidae)	Stem borer	Released 1974- 1985, established	Considerable
	<i>Niphograpta albiputalis</i> Warren (Lepidoptera: Crambidae)	Petiole borer	Released 1990, established	Considerable
	<i>Orthogalumna terebrantis</i> Wallwork (Acari: Sarcoptiformes: Galumnidae)	Leaf miner	Released 1989, established	Considerable
PROTEACEAE				
<i>Hakea gibbosa</i> (Sm.) Cav. (rock hakea) Australia Negligible	<i>Aphanasiumpunctatum</i> (Boisduval) (Coleoptera: Cerambycidae)	Stem borer	Alternative host, released 2001, established	Unknown
	<i>Eryttena consputa</i> Pascoe (Coleoptera: Curculionidae)	Green-seed feeder	Alternative host, released 1975, established	Trivial
<i>Hakea sericea</i> Schrad. & J.C.Wendl. (silky hakea) Australia Substantial	<i>Aphanasiumpunctatum</i> (Boisduval) (Coleoptera: Cerambycidae)	Stem borer	Released 2001, established	Unknown
	<i>Carposina autologa</i> Meyrick (Lepidoptera: Carposinidae)	Seed feeder	Released 1970 & 1982, established	Considerable
	<i>Colletotrichum acutatum</i> J.H. Simmonds f.sp. <i>hakeae</i> Lubbe, Denman, P.F. Cannon, J.Z. Groenew., Lampr. & Crous (Order Incertae sedis: Glomerellaceae)	Stem gummosis disease	Not released, occurs locally; developed as mycoherbicide "Hakatak"	
	<i>Cydiota binotata</i> Lea (Coleoptera: Curculionidae)	Leaf & shoot borer	Released 1979, established	Trivial
	<i>Dicomada rufa</i> Blackburn (Coleoptera: Curculionidae)	Flowerbud feeder	Released 2006, established	Unknown
	<i>Eryttena consputa</i> Pascoe (Coleoptera: Curculionidae)	Green-seed feeder	Released 1970 & 1974, established	Extensive
SALVINIACEAE				
<i>Salvinia molesta</i> D.S.Mitch.	<i>Cyrtobagous salviniae</i> Calder &	Stem borer	Released 1985,	Extensive

(water fern) South America Complete	Sands (Coleoptera: Curculionidae)		established	
SOLANACEAE				
<i>Solanum elaeagnifolium</i> Cav. (silverleaf nightshade/ satansbos) North, Central & South America Substantial	<i>Frumenta nephelomicta</i> (Meyrick) (Lepidoptera: Gelechiidae)	Fruit galler	Released 1979 & 1985, not established	
	<i>Frumenta</i> sp. (Lepidoptera: Gelechiidae)	Stem & fruit galler	Released 1989, not established	
	<i>Leptinotarsa defecta</i> (Stål) (Coleoptera: Chrysomelidae: Chrysomelinae)	Leaf feeder	Released 1992, established	Moderate
	<i>Leptinotarsa texana</i> (Schaeffer) (Coleoptera: Chrysomelidae: Chrysomelinae)	Leaf feeder	Released 1992, established	Extensive
<i>Solanum mauritianum</i> Scop. (bugweed) South America Negligible	<i>Anthonomus santacruzi</i> Hustache (Coleoptera: Curculionidae)	Flowerbud feeder	Released 2008, established	Unknown
	<i>Gargaphia decoris</i> Drake (Hemiptera: Tingidae)	Leaf sucker	Released 1999, established	Trivial
<i>Solanum sisymbriifolium</i> Lam. (wild tomato/ dense- thorned bitter apple) South America Substantial	<i>Gratiana spadicea</i> (Klug) (Coleoptera: Chrysomelidae: Cassidinae)	Leaf feeder	Released 1994, established	Extensive
VERBENACEAE				
<i>Lantana camara</i> L. (lantana) Central & South America Negligible to Substantial (depending on plant variety)	<i>Aceria lantanae</i> (Cook) (Acari: Trombidiformes: Eriophyidae)	Flower galler	Released 2007, established	Extensive, some vars. Coastal
	<i>Alagoasa parana</i> Samuelson (Coleoptera: Chrysomelidae: Galerucinae)	Leaf miner	Released 1985, not established	
	<i>Autoplusia illustrata</i> (Guenée) (Lepidoptera: Noctuidae)	Leaf feeder	Released 1978 & 1984, not established	
	<i>Calycomyza lantanae</i> (Frick) (Diptera: Agromyzidae)	Leaf miner	Released 1982, established	Moderate
	<i>Coelocephalapion camarae</i> Kissinger (Coleoptera: Brentidae: Apioninae)	Petiole galler	Released 2007, established	Unknown
	<i>Crocidosema lantana</i> Busck [formerly in <i>Epinotia</i>] (Lepidoptera: Tortricidae)	Flower- & receptacle miner	Released pre- 1961 & 1984, established	Trivial
	<i>Eutreta xanthochoeta</i> Aldrich (Diptera: Tephritidae)	Shoot galler	Released 1983; not established; rejected	
	<i>Falconia intermedia</i> (Distant) (Hemiptera: Miridae)	Leaf sucker	Released 1999, established	Moderate, localized
	<i>Hypena laceratalis</i> Walker (Lepidoptera: Noctuidae)	Leaf feeder	Not released, native	Moderate
	<i>Lantanophaga pusillidactyla</i> (Walker) (Lepidoptera: Pterophoridae)	Flower miner	Released pre- 1961, established	Trivial
	<i>Leptobrysa decora</i> Drake (Hemiptera: Tingidae)	Leaf sucker	Released 1972- 1984, not established	
	<i>Longitarsus bethae</i> Savini & Escalona (Coleoptera: Chrysomelidae: Galerucinae)	Root feeder	Released 2007, established	Unknown
	<i>Neogalia sunia</i> (Guenée) (Lepidoptera: Noctuidae)	Leaf feeder	Released 1962 & 1969, not established	
	<i>Octotoma championi</i> Baly	Leaf miner	Released 1978,	

	(Coleoptera: Chrysomelidae: Cassidinae)		not established	
	<i>Octotoma scabripennis</i> Guérin-Mèneville (Coleoptera: Chrysomelidae: Cassidinae)	Leaf miner	Released 1971-1975, established	Considerable
	<i>Ophiomyia camarae</i> Spencer (Diptera: Agromyzidae)	Leaf miner	Released 2001, established	Considerable
	<i>Ophiomyia lantanae</i> (Froggatt) (Diptera: Agromyzidae)	Seed miner	Released pre-1961, established	Moderate
	<i>Orthonama ignifera</i> (Warren) [formerly in <i>Leptostales</i>] (Lepidoptera: Geometridae)	Leaf feeder	Approved for release	
	<i>Passalora lantanae</i> (Chupp) U.Braun & Crous var. <i>lantanae</i> [formerly in <i>Mycovellosiella</i>] (anamorphic fungus; Mycosphaerellales: Mycosphaerellaceae)	Leaf pathogen	Released 2002, not established	
	<i>Plagiohammus spinipennis</i> (Thomson) (Coleoptera: Cerambycidae)	Stem borer	Released 1973, not established	
	<i>Salbia haemorrhoidalis</i> (Guenée) (Lepidoptera: Crambidae)	Leaf feeder	Released 1962, established	Trivial
	<i>Teleonemia elata</i> Drake (Hemiptera: Tingidae)	Leaf sucker	Released 1972, not established	
	<i>Teleonemia scrupulosa</i> Stål (Hemiptera: Tingidae)	Leaf & flower sucker	Released 1972, established	Considerable
	<i>Uroplata fulvopustulata</i> Baly (Coleoptera: Chrysomelidae: Cassidinae)	Leaf miner	Released 1978, not established	
	<i>Uroplata girardi</i> Pic (Coleoptera: Chrysomelidae: Cassidinae)	Leaf miner	Released 1974 & 1983, established	Considerable, coastal
	<i>Uroplata lantanae</i> Buzzi & Winder (Coleoptera: Chrysomelidae: Cassidinae)	Leaf miner	Released 1984, not established	

For each of the agents that has established successfully, a rating is given for the perceived degree of damage that it inflicts on the target weed. These ratings (Anon. 1999), which have been widely adopted since 1999, and slightly amended in this account, are sometimes based on formal measurements and otherwise on subjective visual assessments, and include the following categories. *Extensive*: Very high levels of damage, as much as could be expected from the agent. Few plants survive, or growth is arrested, or almost no seeds are produced. *Considerable*: High levels of damage. Some plants may survive but growth rates are noticeably slower, or seed production is reduced by more than 50%. *Moderate*: Perceivable damage, but most plants survive. Growth may be slowed to some extent, or seed production is reduced by less than 50%. *Trivial*: Some damage, but survival, growth and seed production of the plants is almost normal. *Unknown*: Agent has been too recently released, or has not been evaluated.

For each weed species, the combined impact of all agents (i.e. the degree of control) (adapted from Anon. 1999) is rated by estimating the degree to which the impact or importance of the target

weed species has been reduced by the agents. Assessments are based on the degree of reduction in the use of alternative control methods (chemical or mechanical) since the introduction of biological control agents, and include the following categories. *Complete*: No other control measures are needed to reduce the weed to acceptable levels, at least in areas where the agents are established. *Substantial*: Other methods are needed to reduce the weed to acceptable levels, but less effort is required (e.g. less frequent herbicide applications or less herbicide needed per unit area). *Negligible*: In spite of damage inflicted by the agents, control of the weed remains entirely reliant on the implementation of other control measures. *Not determined*: Either the release of the agents has been too recent for meaningful evaluation, or the programme has not been evaluated.

Adapted from: Klein, H. 2011. A catalogue of the insects, mites and pathogens that have been used or rejected, or are under consideration, for the biological control of invasive alien plants in South Africa. In: Moran, V.C., Hoffmann, J.H. & Hill, M.P. (Eds) Biological Control of Invasive Alien Plants in South Africa (1999-2010). *African Entomology* (Special Issue) **19**(2): 515-549.